

PAPER • OPEN ACCESS

## Probing anion resonances in $\text{FeO}^-$ : a species of astrophysical relevance

To cite this article: Roby Chacko *et al* 2017 *J. Phys.: Conf. Ser.* **875** 102019

View the [article online](#) for updates and enhancements.

### Related content

- [Photoelectron imaging of interstellar medium anions](#)  
Nrisimhamurty Madugula, Roby Chacko, Pranawa C Deshmukh *et al.*
- [COLLISIONAL DESTRUCTION OF  \$\text{FeC}^{\(n\)-}\$  \( \$n = 1\$  TO 4, 6\) ANIONS OF ASTROPHYSICAL RELEVANCE](#)  
M. Nrisimhamurty, R. G. Mane, Roby Chacko *et al.*
- [Astrophysical relevance of storage-ring electron-ion recombination experiments](#)  
Stefan Schippers



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

## Probing anion resonances in $\text{FeO}^-$ : a species of astrophysical relevance

Roby Chacko<sup>†1</sup>, Shreyak Banhatti<sup>†2</sup>, A. K. Gupta<sup>\*3</sup>, and G. Aravind<sup>†4</sup>

<sup>†</sup> Department of Physics, Indian Institute of Technology Madras, Chennai - 600 036, India


\* Nuclear Physics Division, Bhabha Atomic Research Centre, Mumbai 400 085, India

**Synopsis** We report the first experimental probe on resonance states in  $\text{FeO}^-$  anion, a molecule of astrophysical relevance. Excitation of  $\text{FeO}^-$  through collision with argon was performed to measure the kinetic energy release in the centre of mass frame. Results show access of two anion resonances upon collisional excitation. We discuss our evaluation of these two resonances from kinetic energy release in the centre of mass frame. The astrophysical importance of these anion resonances will be presented.

$\text{FeO}$  is the first iron-bearing molecule discovered in the inter-stellar medium (ISM) [1], [2]. Depletion of refractory elements in the ISM motivates the search and study of iron-bearing ISM molecules. Recently, another iron-bearing molecule,  $\text{FeCN}$  was unambiguously identified in the envelope of carbon-rich asymptotic giant branch star, IRC+10216 [3]. Although ISM anions were predicted in 1981 by Herbst *et al.* [4], only in 2006 the first ISM anion  $\text{C}_6\text{H}^-$  was detected [5]. This was followed by recent detections of  $\text{CN}^-$ ,  $\text{C}_3\text{N}^-$ ,  $\text{C}_4\text{H}^-$ ,  $\text{C}_6\text{H}^-$ ,  $\text{C}_8\text{H}^-$  and  $\text{C}_5\text{N}^-$ . Anion resonances are states embedded in the detachment continuum and are short-lived states. They are formidable to study theoretically due to high electron correlation. Hence experimental investigations such as dissociative electron attachment [6], collisional or photoexcitation become important to probe these resonances. In the present work, we have probed  $\text{FeO}^-$  which is detected in its neutral state in the ISM. Resonance states of  $\text{FeO}^-$  can be accessed by electron attachment to  $\text{FeO}$  in the ISM and would result in autodetachment or dissociation. Anion resonances thus critically determine the ISM reactions and relative abundances of various ISM species.

$\text{FeO}^-$  were produced in Cs-sputter source and were accelerated to 15keV and eventually collided with Argon in a collision cell. The collisionally excited anion dissociates and the anionic daughters scattered in the forward and backward directions were only detected using an energy analyser. Two daughter anions were detected in our experiments as shown in the figure. The kinetic energy (KE) released in the center of mass frame was evaluated from the lab frame KE release. The energy released in the center of mass frame for both the daughter anions were

different, indicating access of two different anionic states in this experiment. In our recent work on  $\text{FeC}^-$  anions we have observed two anion resonances [7]. We have evaluated the ground state of the  $\text{FeO}^-$  and its vibrational wavefunctions using LEVEL program [7]. We employ reflection method, on the CM frame energy release to deduce the two anion resonances accessed in our experiment. Our results are of paramount importance in studying the dissociation dynamics involving iron containing ISM molecules. The important implications of our results to astronomy will be discussed in the presentation.



**Figure 1.** Daughter-ion peaks recorded for the CID of  $\text{FeC}^-$  anion

### References

- [1] C. M. Walmsley *et al.* 2002 *Astrophys. J.* **566** L109
- [2] R. S. Furuya *et al.* 2003 *Astr. Astrophys.* **409** L21
- [3] L. N. Zack *et al.* 2011 *Astrophys. J. Lett.* **733** L36
- [4] E. Herbst 1981 *Nature* **289** 656
- [5] M. C. McCarthy *et al.* 2006 *Astrophys. J. Lett.* **652** L141
- [6] V. S. Prabhudesai *et al.* 2005 *Phys. Rev. Lett.* **95** 143202
- [7] M. Nrisimhamurty *et al.* 2016 *Astrophys. J.* **833** 269

<sup>1</sup>E-mail: [robychako@gmail.com](mailto:robychako@gmail.com)

<sup>2</sup>E-mail: [shreyak.rocker@gmail.com](mailto:shreyak.rocker@gmail.com)

<sup>3</sup>E-mail: [anit@tifr.res.in](mailto:anit@tifr.res.in)

<sup>4</sup>E-mail: [garavind@iitm.ac.in](mailto:garavind@iitm.ac.in)

