SZEGÖ AND WIDOM THEOREMS FOR THE NEIL ALGEBRA

SRIRAM BALASUBRAMANIAN¹, SCOTT MCCULLOUGH², AND UDENI WIJESOORIYA

In appreciation for his profound influence on operator theory and our mathematical lives, we dedicate this article to Joe Ball.

ABSTRACT. Versions of well known function theoretic operator theory results of Szegö and Widom are established for the Neil algebra. The Neil algebra is the subalgebra of the algebra of bounded analytic functions on the unit disc consisting of those functions whose derivative vanishes at the origin.

1. INTRODUCTION

Let \mathbb{C} denote the complex numbers, $\mathbb{D} = \{|z| < 1\} \subseteq \mathbb{C}$ denote the unit disk with its boundary $\mathbb{T} = \{|z| = 1\}$. Denote by $H^2 = H^2(\mathbb{D})$ and $H^\infty = H^\infty(\mathbb{D})$ the standard Hardy spaces of functions analytic in \mathbb{D} with square summable power series coefficients and bounded analytic functions on \mathbb{D} respectively. Let L^p denote the L^p spaces for the \mathbb{T} (identified with the corresponding L^p spaces for $[0, 2\pi]$ with respect to the measure $\frac{dt}{2\pi}$). Let \mathscr{P} denote the set of analytic polynomials that vanish at 0. Thus a $p \in \mathscr{P}$ has the form,

$$p(z) = \sum_{j=1}^{n} p_j z^j$$

for some positive integer n and $p_1, \ldots, p_n \in \mathbb{C}$. Given a non-negative function ρ on \mathbb{T} with $\log(\rho) \in L^1$ a (special case of a) well known result of Szegö (see for instance [14] page 219) identifies the $L^2(\rho)$ distance from the constant function 1 to \mathscr{P} .

Theorem 1.1 (of Szegö).

$$\inf\{\int_0^{2\pi} |p-1|^2 \, \rho \, \frac{dt}{2\pi} : p \in \mathscr{P}\} = \exp(\int_0^{2\pi} \log(\rho) \, \frac{dt}{2\pi}).$$

A theorem of Widom characterizes those unimodular functions $\phi \in L^{\infty}$ whose distance to H^{∞} is less than one in terms of Toeplitz operators. A $\phi \in L^{\infty}$ induces a

¹⁹⁹¹ Mathematics Subject Classification. 47B335, 30H10 (Primary) 30H05, 46E20 (secondary).

Key words and phrases. Toeplitz operators, Szegö's Theorem, constrained algebra, Neil algebra, distinguished variety.

¹Supported by the New Faculty Initiative Grant (MAT/15-16/836/NFIG/SRIM) of IIT Madras. ²Research supported by the NSF grant DMS-1361501.

multiplication operator $M_{\phi}: L^2 \to L^2$ defined by $M_{\phi}f = \phi f$. Let $V: H^2 \to L^2$ denote the inclusion. The operator $T_{\phi} = V^* M_{\phi} V$ is the *Toeplitz operator* with symbol ϕ .

Theorem 1.2 (Widom's invertibility criteria [11, Theorem 7.30]). Suppose $\phi \in L^{\infty}$ is unimodular. There exists an $f \in H^{\infty}$ such that $||f - \phi|| < 1$ if and only if T_{ϕ} is left invertible.

Sarason [18] established a version of Theorem 1.1 for the annulus and Abrahamse [1, Theorems 4.1 and 4.6] established a version of Theorem 1.2 for multiply connected domains. In this paper we establish Szegö and Widom type theorems for the *Neil* algebra. The Neil algebra \mathfrak{A} is the subalgebra of $H^{\infty}(\mathbb{D})$ consisting of those functions whose derivative vanishes at 0. It is perhaps the simplest example of a constrained algebra. As with extending classical results from the unit disc to multiply connected domains, here it is necessary to replace H^2 with a family of Hilbert-Hardy spaces that parameterize the distinction between harmonic functions and the real parts of analytic functions in \mathfrak{A} either explicitly or implicitly in the statement of the results and their proofs. In addition to the references already cited, see for instance [2, 3, 16, 8] for related results on multiply connected domains, [5, 6, 7, 12, 10, 16, 17] for results on constrained algebras, [4] for results in the context of uniform algebras and finally [13] for a Pick interpolation theorem on distinguished varieties. Let \mathfrak{A}_0 denote those functions in \mathfrak{A} that vanish at 0. Hence $\mathfrak{A}_0 = z^2 H^{\infty}$.

Theorem 1.3 (Szegö Theorem for \mathfrak{A}). ¹ Suppose $\rho > 0$ is a continuous function on \mathbb{T} and let

$$C_{\rho} = \int_0^{2\pi} \log(\rho) \, \frac{dt}{2\pi}, \quad \lambda = \int_0^{2\pi} \rho(t) \exp(-it) \, \frac{dt}{2\pi}$$

With these notations,

$$\inf\{\int_0^{2\pi} |1-p|^2 \rho \frac{dt}{2\pi} : p \in \mathfrak{A}_0\} = \exp(C_\rho) + \exp(-C_\rho) |\lambda|^2.$$

Remark 1.4. Note that $\lambda = 0$ if and only if 1 and e^{it} are orthogonal in $L^2(\rho)$ and in this case it is evident that the distance from 1 to \mathscr{P} is the same as the distance from 1 to the subspace \mathfrak{A}_0 of \mathscr{P} .

To state the analog of Theorem 1.2 for \mathfrak{A} some notations are needed. Let $\mathbb{B}^2 = \{(z,w) \in \mathbb{C}^2 : |z^2| + |w|^2 = 1\}$ denote the unit ball in \mathbb{C}^2 . To $\alpha = (a,b) \in \mathbb{B}^2$ associate the subspace $H^2_{\alpha} \subseteq H^2$ consisting of those $f \in H^2$ such that

$$f(0) b = f'(0) a.$$

¹[4, Theorem 5.1] covers the case $\lambda = 0$.

Let $V_{\alpha}: H^2_{\alpha} \to L^2$ denote the inclusion. Hence $P_{\alpha} = V_{\alpha}V^*_{\alpha}: L^2 \to H^2_{\alpha}$ is the projection onto H^2_{α} . Given $\phi \in L^{\infty}$, define $T^{\alpha}_{\phi}: H^2_{\alpha} \to H^2_{\alpha}$ by

$$T^{\alpha}_{\phi} = V^*_{\alpha} M_{\phi} V_{\alpha}.$$

It is the Toeplitz operator with symbol ϕ with respect to α [8]. In particular, if $\phi \in \mathfrak{A}$ and $f \in H^2_{\alpha}$, then $V^*T^{\alpha}_{\phi}f = \phi f = T^{\alpha}_{\phi}f$.

Remark 1.5. Given $\alpha = (a, b)$ and $\beta = (c, d)$, if ad = bc, then $H^2_{\alpha} = H^2_{\beta}$ and likewise $T^{\alpha}_{\phi} = T^{\beta}_{\phi}$. Thus, \mathbb{P} , complex projective space obtained by moding out \mathbb{B}^2 by the relation (a, b) = (c, d), is a natural choice of parameter space. For ease of exposition we accept the redundancy inherent in the use of \mathbb{B}^2 .

Theorem 1.6 (Inversion for \mathfrak{A}). Suppose $\phi \in L^{\infty}$ is unimodular. The distance from ϕ to \mathfrak{A} is strictly less than one if and only if T^{α}_{ϕ} is left invertible for each $\alpha \in \mathbb{B}^2$. Likewise, the distance from ϕ to the invertible elements of \mathfrak{A} is strictly less than one if and only if T^{α}_{ϕ} is invertible for each $\alpha \in \mathbb{B}^2$.

Before turning to the proofs of Theorems 1.3 and 1.6, we pause to introduce some conventions and basic background on the spaces H^2_{α} . For $p = 2, \infty$, the standard identification of $H^p(\mathbb{D})$ with $H^p(\mathbb{T})$, where the latter is viewed as the subspace of $L^p(\mathbb{T})$ consisting of those f with vanishing negative Fourier coefficients, will be used routinely and without comment. Let H^2_1 denote the subspace of H^2 consisting of those $f \in H^2$ whose Fourier coefficient

$$\hat{f}(1) = \int_0^{2\pi} f \, e^{-it} \, \frac{dt}{2\pi} = 0.$$

Evidently, H_1^2 is the closure of \mathfrak{A} in H^2 . The following Lemma can be found in [10] for instance. The first part follows from the easily verified fact that $\{a + bz, z^n : n \geq 2\}$ is an orthonormal basis for H_{α}^2 ; and the moreover part, from a standard reproducing kernel Hilbert space argument.

Lemma 1.7. For each $\alpha = (a, b) \in \mathbb{B}^2$, the space H^2_{α} has reproducing kernel,

$$k_w^{\alpha}(z) = k^{\alpha}(z, w) = (a + bz)\overline{(a + bw)} + \frac{z^2 \overline{w^2}}{1 - z\overline{w}}, \quad z, w \in \mathbb{D}.$$

In particular,

$$||k_0^{\alpha}||^2 = k^{\alpha}(0,0) = |a|^2,$$

and thus $k_w^{\alpha} \neq 0$ with the exception of $\alpha = (0, 1)$ and w = 0.

Moreover, if $\psi \in \mathfrak{A}$ and $w \in \mathbb{D}$, then $(T_{\psi}^{\alpha})^* k_w^{\alpha} = \overline{\psi(w)} k_w^{\alpha}$.

2. Proof of Theorem 1.3

As a first step, observe that it suffices to prove the theorem under the additional hypothesis that $C_{\rho} = 0$. Indeed, if not, let $\tilde{\rho} = \exp(-C_{\rho})\rho$, so that $\int_{0}^{2\pi} \log(\tilde{\rho}) \frac{dt}{2\pi} = 0$. In particular, $C_{\tilde{\rho}} = 0$ and with

$$\tilde{\lambda} = \int_0^{2\pi} \tilde{\rho} \, \exp(-it) \, \frac{dt}{2\pi} = \exp(-C_{\rho}) \, \lambda,$$

if Theorem 1.3 holds for $\tilde{\rho}$, then

$$\inf\{\int_0^{2\pi} |p-1|^2 \,\tilde{\rho} \,\frac{dt}{2\pi} : p \in \mathfrak{A}_0\} = 1 + |\tilde{\lambda}|^2.$$

Thus,

$$\inf\{\int_{0}^{2\pi} |p-1|^{2} \rho \frac{dt}{2\pi} : p \in \mathfrak{A}_{0}\} = \exp(C_{\rho}) \inf\{\int_{0}^{2\pi} |p-1|^{2} \tilde{\rho} dtn : p \in \mathfrak{A}_{0}\}$$
$$= \exp(C_{\rho})(1+|\tilde{\lambda}|^{2}) = \exp(C_{\rho}) + \exp(-C_{\rho}) |\lambda|^{2}$$

as claimed. Accordingly, for the remainder of the proof, assume $C_{\rho} = 0$.

Let

$$\sigma = \frac{1}{\sqrt{1+|\lambda|^2}} (1,\lambda) \in \mathbb{B}^2$$

In particular,

$$\|k_0^{\sigma}\|^2 = \frac{1}{1+|\lambda|^2}.$$

Note that, as sets, $L^2(\rho)$ and L^2 are the same and thus we may consider H^2 as a Hilbert space with the alternate inner product,

$$\langle f,g\rangle_{\rho} = \int_0^{2\pi} f\overline{g}\,\rho \frac{dt}{2\pi}.$$

To keep the distinction clear, denote this latter space by $H^2(\rho)$. Since the closure of \mathfrak{A}_0 in $H^2(\rho)$ is $z^2H^2 = z^2H^2(\rho)$, the objective is to find the $H^2(\rho)$ -distance from 1 to z^2H^2 . That is, to show

$$\inf\{\int_0^{2\pi} |p-1|^2 \,\tilde{\rho} \,\frac{dt}{2\pi} : f \in z^2 H^2\} = 1 + |\tilde{\lambda}|^2.$$

Since ρ is continuous and strictly positive, $\log(\rho)$ is continuous. It has Fourier series expansion

$$\log(\rho) = \sum_{j=-\infty}^{\infty} c_j e^{ijt},$$

where, because it is real-valued, $c_{-j} = \overline{c_j}$. Moreover, $c_0 = 0$ and $c_1 = \lambda$, since $C_{\rho} = 0$ and by the very definitions of C_{ρ} and λ . Letting γ denote the H^2 function represented by the series

$$\gamma = \sum_{j=1}^{\infty} c_j e^{ijt},$$

it follows that $\log(\rho) = \gamma + \gamma^*$ as elements of L^2 . Further, since

$$|\exp(\pm\gamma)|^2 = \exp\left(\pm(\gamma+\gamma^*)\right) = \rho^{\pm 1},$$

both $\exp(\pm\gamma)$ are in H^{∞} . The mapping $U: H^2(\rho) \to H^2$ defined by $Uf = \exp(\gamma)f$ is a unitary map with inverse $U^*f = \exp(-\gamma)f$. Moreover, $U(z^2H^2) = z^2H^2$. Thus, the aim is to find the H^2 -distance from $\exp(\gamma)$ to z^2H^2 .

Given $f \in z^2 H^2$, let $g = \exp(\gamma) - f$ and estimate, using g(0) = 1 and the Cauchy-Schwarz inequality,

(2.1)
$$\begin{aligned} \|\exp(\gamma) - f\|^{2} &= \|g\|^{2} \\ &\geq \frac{|\langle g, k_{0}^{\sigma} \rangle|^{2}}{\|k_{0}^{\sigma}\|^{2}} \\ &= |g(0)|^{2} \left(1 + |\lambda|^{2}\right) \\ &= 1 + |\lambda|^{2}. \end{aligned}$$

Let

$$f = \exp(\gamma) - (1 + |\lambda|^2)k_0^{\sigma}$$

and note f(0) = 0 and $f'(0) = \gamma'(0) - \lambda = 0$. Thus $f \in z^2 H^2$ and, with this choice of f, equality holds in the Cauchy-Schwarz inequality in equation (2.1).

3. Toeplitz operators on \mathfrak{A}

This section contains the proof of Theorem 1.6.

Lemma 3.1. If $\phi \in L^{\infty}$, then $||T^{\alpha}_{\phi}|| = ||\phi||$ and $(T^{\alpha}_{\phi})^* = T^{\alpha}_{\overline{\phi}}$.

Proof. Since $M_{\phi}^* = M_{\overline{\phi}}$, it follows that $(T_{\phi}^{\alpha})^* = V_{\alpha}^* M_{\phi}^* V_{\alpha} = V_{\alpha}^* M_{\overline{\phi}} V_{\alpha} = T_{\overline{\phi}}^{\alpha}$. Since V_{α} is an isometry, it follows that $||T_{\phi}^{\alpha}|| \leq ||M_{\phi}|| = ||\phi||$. Now let $V : H^2 \to L^2$ and $W : z^2 H^2 \to L^2$ denote the inclusion maps. In particular, $V^* M_{\phi} V$ is T_{ϕ} , the usual Toeplitz operator with symbol ϕ . On the other hand, $W^* M_{\phi} W = W^* T_{\phi}^{\alpha} W$. With $U : H^2 \to z^2 H^2$ given by $Uf = z^2 f$, it follows that U is unitary and, for $f, g \in H^2$,

$$\langle M_{\phi}WUf, WUg \rangle = \langle z^{2}\phi f, z^{2}g \rangle = \langle \phi f, g \rangle = \langle M_{\phi}f, g \rangle = \langle V^{*}M_{\phi}Vf, g \rangle$$

Hence $U^*W^*M_{\phi}WU = V^*M_{\phi}V = T_{\phi}$ and consequently $W^*T^{\alpha}_{\phi}W$ is unitarily equivalent to T_{ϕ} . Hence $||T^{\alpha}_{\phi}|| \ge ||T_{\phi}||$. Since, as is well known that $||T_{\phi}|| = ||\phi||$ ([15]), the result follows.

Let $\mathscr{B}(L^2)$ denote the bounded linear operators on L^2 .

Lemma 3.2. Giving \mathbb{B}^2 its usual topology and $\mathscr{B}(L^2)$ its norm topology, the mapping $\mathbb{B}^2 \ni \alpha \to P_\alpha \in \mathscr{B}(L^2)$ is continuous.

Proof. Since $\{a + bz, z^n : n \ge 2\}$ is an orthonormal basis for H^2_{α} , if $f = \sum f_n z^n \in H^2$ and $\alpha = (a, b) \in \mathbb{B}^2$, then

$$P_{\alpha}f = (\overline{a}f_0 + \overline{b}f_1)(a + bz) + \sum_{n=2}^{\infty} f_n z^n.$$

Thus, letting Q denote the projection onto $z^2 H^2$ and $F_{\alpha} = (a + bz)$ (a unit vector),

$$P_{\alpha} = F_{\alpha}F_{\alpha}^* + Q$$

where $F_{\alpha}F_{\alpha}^{*}: L^{2} \to L^{2}$ is the rank one projection operator,

$$F_{\alpha}F_{\alpha}^{*}f = \langle f, F_{\alpha}\rangle F_{\alpha} = (\overline{a}f_{0} + bf_{1})F_{\alpha}.$$

Thus, if $\beta = (c, d) \in \mathbb{B}^2$, then

$$P_{\alpha} - P_{\beta} = F_{\alpha}F_{\alpha}^* - F_{\beta}F_{\beta}^* = F_{\alpha}(F_{\alpha} - F_{\beta})^* + (F_{\alpha} - F_{\beta})F_{\beta}^*.$$

Since $||F_{\alpha} - F_{\beta}|| = ||\alpha - \beta||$, the result follows.

Let $\mathcal{M}\subseteq L^1$ denote the subspace consisting of those L^1 functions with Fourier series of the form

(3.1)
$$\hat{f}(-1)\exp(-it) + \sum_{j=1}^{\infty} \hat{f}(j)\exp(ijt).$$

The following lemma is the $\mathscr M$ version of the well known factorization theorem for H^1 functions.

Lemma 3.3. If $h \in \mathcal{M}$, then there exist

```
(i) \alpha \in \mathbb{B}^{2};

(ii) f \in H_{\alpha}^{2}; and

(iii) g \in L^{2}

such that

(a) \overline{g} \in (H_{\alpha}^{2})^{\perp};

(b) h = fg; and

(c) \|h\|_{1} = \|f\|_{2} \|g\|_{2}.
```

Proof. The function $\psi = zh$ is in H^1 and therefore there exists $F, G \in H^2$ such that zh = FG and $\|h\|_1 = \|\psi\|_1 = \|F\|_2 \|G\|_2$ [11, Corollary 6.27]. Moreover, since $\psi'(0) = 0$, it follows that F'(0) G(0) + F(0) G'(0) = 0. There is an $\alpha = (a, b) \in \mathbb{B}^2$ such that $F \in H^2_{\alpha}$. (Indeed, simply choose $\alpha \in \mathbb{B}^2$ such that aF'(0) = bF(0).) Thus there is a constant c and an H^2 function F_0 such that

$$F = c(a+bz) + z^2 F_0.$$

Hence, there is a constant d and H^2 function G_0 such that

$$G = d(a - bz) + z^2 G_0.$$

Let $g = \overline{z}G$, in which case h = Fg and $||g||_2 = ||G||_2$. Moreover,

$$\langle a+bz,\overline{g}\rangle = d \int_0^{2\pi} (a+bz) \left(d(a\overline{z}-b)+zG_0\right) \frac{dt}{2\pi} = 0$$

and, for $n \geq 2$,

$$\langle z^n, \overline{g} \rangle = \int_0^{2\pi} z^n \left(d(a\overline{z} - b) + zG_0 \right) \frac{dt}{2\pi} = 0.$$

Hence $\overline{g} \in (H^2_{\alpha})^{\perp}$.

Recall $(L^1)^* = L^\infty$ with the equality interpreted as the isometric isomorphism determined by the mapping that assigns to $\phi \in L^\infty$ the linear functional $\lambda_\phi : L^1 \to \mathbb{C}$ given by

$$\lambda_{\phi}(\psi) = \int_0^{2\pi} \phi \,\psi \,\frac{dt}{2\pi}.$$

Moreover, letting

$$\mathscr{M}^{\perp} := \{ \phi \in L^{\infty} : \int_{0}^{2\pi} \phi \, \psi \frac{dt}{2\pi} = 0, \text{ for all } \psi \in \mathscr{M} \},$$

and $\pi: L^{\infty} \to L^{\infty}/\mathscr{M}^{\perp}$ denote the quotient mapping, the mapping $\Lambda: L^{\infty}/\mathscr{M}^{\perp} \to \mathscr{M}^*$ given by

$$\Lambda(\pi(\lambda_{\phi})) = (\lambda_{\phi})|_{\mathscr{M}},$$

is an isometric isomorphism. Finally, if $\phi \in \mathcal{M}$ and $\psi \in \mathfrak{A}$, then

$$\int_0^{2\pi} \phi \,\psi \,\frac{dt}{2\pi} = 0.$$

Thus, $\mathfrak{A} \subseteq \mathscr{M}^{\perp}$. On the other hand, $e^{ijt} \in \mathscr{M}$ for $j = -1, 1, 2, \ldots$ and therefore if $\psi \in \mathscr{M}^{\perp}$, then its Fourier series has the form

$$\psi = \hat{\psi}(0) + \sum_{j=2}^{\infty} \hat{\psi}(j) e^{ijt}.$$

Hence $\psi \in \mathfrak{A}$ and thus we may view Λ as having domain L^{∞}/\mathfrak{A} . The following lemma summarizes the discussion (see [9, page 88]).

Lemma 3.4. $\Lambda : L^{\infty}/\mathfrak{A} \to \mathscr{M}^*$ defined by sending $\pi(\phi)$ to the linear functional $\lambda_{\phi} : \mathscr{M} \to \mathbb{C}$ given by

$$\tilde{\lambda}_{\phi}(f) = \int_{0}^{2\pi} \phi f \, \frac{dt}{2\pi}$$

is an isometric isomorphism.

Lemma 3.5. If $\phi \in L^{\infty}$ and $\psi \in \mathfrak{A}$, then

$$T^{\alpha}_{\overline{\psi}\phi} = T^{\alpha}_{\overline{\psi}} T^{\alpha}_{\phi}$$
$$T^{\alpha}_{\overline{\psi}\phi} = T^{\alpha}_{\overline{\phi}} T^{\alpha}_{\psi}.$$

Proof. Let $f, g \in H^2_{\alpha}$ be given. Since $\psi g \in H^2_{\alpha}$, it follows, using Lemma 3.1, that $\langle T^{\alpha}_{\overline{\psi}} T^{\alpha}_{\phi} f, g \rangle = \langle T^{\alpha}_{\phi} f, (T^{\alpha}_{\overline{\psi}})^* g \rangle = \langle T^{\alpha}_{\phi} f, T^{\alpha}_{\psi} g \rangle = \langle V^*_{\alpha} \phi f, \psi g \rangle = \langle \phi f, V_{\alpha} \psi g \rangle = \langle \phi f, \psi g \rangle = \langle \overline{\psi} \phi f, g \rangle = \langle \overline{\psi} \phi V_{\alpha} f, V_{\alpha} g \rangle = \langle T^{\alpha}_{\overline{\psi}} f, g \rangle$. Thus $T^{\alpha}_{\overline{\psi}\phi} = T^{\alpha}_{\overline{\psi}} T^{\alpha}_{\phi}$. Applying Lemma 3.1 to what has already been proved, $T^{\alpha}_{\psi\overline{\phi}} = (T^{\alpha}_{\overline{\psi}\phi})^* = (T^{\alpha}_{\overline{\psi}} T^{\alpha}_{\phi})^* = T^{\alpha}_{\overline{\phi}} T^{\alpha}_{\psi}$.

An element $\psi \in \mathfrak{A}$ is *invertible in* \mathfrak{A} if it does not vanish in \mathbb{D} and $\psi^{-1} = \frac{1}{\psi} \in \mathfrak{A}$.

Lemma 3.6. Suppose $\psi \in \mathfrak{A}$. The following are equivalent.

- (i) ψ is invertible in \mathfrak{A} ;
- (ii) there is an $\alpha \in \mathbb{B}^2$ such that T^{α}_{ψ} is right invertible;
- (iii) T^{α}_{ψ} is invertible for each $\alpha \in \mathbb{B}^2$.

Moreover, in this case $(T^{\alpha}_{\psi})^{-1} = T^{\alpha}_{\psi^{-1}}$.

Proof. Evidently item (i) implies item (ii) implies item (ii). Now suppose there is an α such that $T := T_{\psi}^{\alpha}$ is right invertible. The Hilbert space H_{α}^{2} has a reproducing kernel $k_{w}^{\alpha}(z)$ and further $T^{*}k_{w}^{\alpha} = \overline{\psi(w)}k_{w}^{\alpha}$ by Lemma 1.7. Since T is right invertible, T^{*} is bounded below; i.e., there is a $\delta > 0$ such that $||T^{*}f|| \geq \delta ||f||$ for all $f \in H_{\alpha}^{2}$. Hence,

$$\overline{|\psi(w)|} \|k_w^{\alpha}\| = \|T^*k_w^{\alpha}\| \ge \delta \|k_w^{\alpha}\|.$$

Moreover, by Lemma 1.7 $k_w^{\alpha} \neq 0$ for $w \neq 0$. Thus $|\frac{1}{\psi}(w)| \leq \frac{1}{\delta}$ for $w \in \mathbb{D} \setminus \{0\}$ and therefore, as $\frac{1}{\psi}$ is otherwise analytic, $|\frac{1}{\psi}|$ is bounded by $\frac{1}{\delta}$. Since $\psi \in \mathfrak{A}$ it follows that $\frac{1}{\psi} \in \mathfrak{A}$ too; i.e., item (i) holds.

Lemma 3.7. Suppose $\phi \in L^{\infty}$ is unimodular. If there exists $\psi \in \mathfrak{A}$ such that $\|\phi - \psi\| < 1$, then $T^{\alpha}_{\overline{\phi}}T^{\alpha}_{\psi}$ is invertible, and therefore T^{α}_{ϕ} is left invertible, for each $\alpha \in \mathbb{B}^2$. Further, if ψ is invertible in \mathfrak{A} , then $T^{\alpha}_{\overline{\psi}}T^{\alpha}_{\phi}$ is invertible, and therefore T^{α}_{ϕ} is invertible, for each $\alpha \in \mathbb{B}^2$.

Proof. Suppose there exists $\psi \in \mathfrak{A}$ such that $\|\phi - \psi\| < 1$. In this case $\|1 - \psi\overline{\phi}\| < 1$, since $|\phi| = 1$ (unimodular). Hence, by Lemma 3.1, for a given $\alpha \in \mathbb{B}^2$,

$$1 > \|1 - \psi \overline{\phi}\| = \|T^{\alpha}_{1 - \psi \overline{\phi}}\| = \|1 - T^{\alpha}_{\psi \overline{\phi}}\|$$

In particular, $T^{\alpha}_{\psi\phi}$ is invertible. Since $\psi \in \mathfrak{A}$, Lemma 3.5 applies to give, $T^{\alpha}_{\psi\phi} = T^{\alpha}_{\phi}T^{\alpha}_{\psi}$. Thus T^{α}_{ϕ} is right invertible. By Lemma 3.1, $(T^{\alpha}_{\phi})^* = T^{\alpha}_{\phi}$ is left invertible.

Now, assuming ψ is invertible in \mathfrak{A} , by Lemma 3.6, T_{ψ}^{α} is invertible. The invertibility of T_{ϕ}^{α} follows. Thus, again using Lemma 3.1, T_{ϕ}^{α} is invertible.

Lemma 3.8. If $\phi \in L^{\infty}$ and T^{α}_{ϕ} is left invertible for each $\alpha \in \mathbb{B}^2$, then there exists an $\epsilon \in (0, 1]$, such that for each $\alpha \in \mathbb{B}^2$ and $f \in H^2_{\alpha}$,

$$||T^{\alpha}_{\phi}f|| \ge \epsilon ||f||.$$

Proof. For $\alpha \in \mathbb{B}^2$, define $X_{\alpha} : L^2 \to L^2$ by $X_{\alpha} = P_{\alpha}M_{\phi}P_{\alpha} + (I-P_{\alpha})$. Given $\alpha \in \mathbb{B}^2$, since T_{ϕ}^{α} is left invertible, there exists an $\epsilon_{\alpha} \in (0, 1]$ such that $\|V_{\alpha}T_{\phi}^{\alpha}f\| = \|T_{\phi}^{\alpha}f\| \ge \epsilon_{\alpha}\|f\|$ for $f \in H_{\alpha}^2$. Hence, given F = f + g with $f \in H_{\alpha}^2$ and $g \in (H_{\alpha}^2)^{\perp}$,

$$||X_{\alpha}F||^{2} = ||V_{\alpha}T_{\phi}^{\alpha}f||^{2} + ||g||^{2} \ge \epsilon_{\alpha}^{2}||F||^{2}$$

Thus, $||X_{\alpha}F|| \ge \epsilon_{\alpha}||F||$ for all $F \in L^2$.

To show there is an $\epsilon > 0$ such that $||X_{\alpha}F|| \ge \epsilon ||F||$ for all $\alpha \in \mathbb{B}^2$ and $F \in L^2$, we argue by contradiction. Accordingly suppose no such $\epsilon > 0$ exists. By compactness of \mathbb{B}^2 , there is a sequence $\alpha_n = (a_n, b_n)$ from \mathbb{B}^2 , that, by passing to a subsequence if needed, we may assume converges to some $\beta = (a, b) \in \mathbb{B}^2$ and a unit vectors $F_n \in L^2$ such that $(||X_{\alpha_n}F_n||)_n$ converges to 0. But then,

$$0 < \epsilon_{\beta} \le \|X_{\beta}F_n\| \le \|X_{\alpha_n}F_n\| + \|(X_{\beta} - X_{\alpha_n})F_n\|.$$

By norm continuity (Lemma 3.2) the last term on the right hand side tends to 0 and by assumption the first term on the right hand side tends to 0, a contradiction.

To complete the proof, simply observe if $f \in H^2_{\alpha} \subseteq L^2$, then $\|\phi\| \|f\| \ge \|T^{\alpha}_{\phi}f\| = \|X_{\alpha}f\| \ge \epsilon \|f\|$.

Lemma 3.9. Suppose $\phi \in L^{\infty}$ is unimodular. The distance from ϕ to \mathfrak{A} is strictly less than one if and only if T^{α}_{ϕ} is left invertible for each $\alpha \in \mathbb{B}^2$.

Proof. Suppose T^{α}_{ϕ} is left invertible for each $\alpha \in \mathbb{B}^2$. In this case, Lemma 3.8 applies and thus there is an $1 \ge \epsilon > 0$ such that for each α and $f \in H^2_{\alpha}$,

$$\|T^{\alpha}_{\phi}f\| \ge \epsilon \|f\|.$$

Now let $h \in \mathscr{M}$ be given. By Lemma 3.3 there is an $\alpha \in \mathbb{B}^2$ and $f \in H^2_{\alpha}$ and a $g \in L^2$ such that $\overline{g} \in (H^2_{\alpha})^{\perp}$ and both h = fg and $\|h\|_1 = \|f\|_2 \|g\|_2$. Thus,

$$\int_{0}^{2\pi} \phi h \frac{dt}{2\pi} \bigg| = \bigg| \int_{0}^{2\pi} \phi fg \frac{dt}{2\pi} \bigg|$$
$$= |\langle \phi f, \overline{g} \rangle|$$
$$= |\langle \phi f, (I - P_{\alpha})\overline{g} \rangle|$$
$$= |\langle (I - P_{\alpha})\phi f, \overline{g} \rangle|$$
$$\leq ||(I - P_{\alpha})\phi f|| ||g||.$$

On the other hand, using the unimodular hypothesis,

$$\begin{split} \|f\|^{2} &= \|\phi f\|^{2} = \|P_{\alpha}\phi f\|^{2} + \|(I - P_{\alpha})\phi f\|^{2} \\ &= \|T_{\alpha}^{\phi}f\|^{2} + \|(I - P_{\alpha})\phi f\|^{2} \\ &\geq \epsilon^{2}\|f\|^{2} + \|(I - P_{\alpha})\phi f\|^{2}. \end{split}$$

Thus, $(1 - \epsilon^{2})\|f\|^{2} \geq \|(I - P_{\alpha})\phi f\|^{2}$. Therefore,
 $\left|\int_{0}^{2\pi} \phi h \frac{dt}{2\pi}\right| \leq \sqrt{1 - \epsilon^{2}} \|f\|_{2} \|g\|_{2} = \sqrt{1 - \epsilon^{2}} \|h\|_{1}. \end{split}$

By Lemma 3.4, it now follows that $\|\pi(\phi)\| < 1$, where $\pi : L^{\infty} \to L^{\infty}/\mathfrak{A}$ is the quotient map; i.e., the distance from ϕ to \mathfrak{A} is less than one.

Conversely, if the distance from ϕ to \mathfrak{A} is less than one, then there exists a $\psi \in \mathfrak{A}$ such that $\|\phi - \psi\| < 1$. It follows from Lemma 3.7 that T_{ϕ}^{α} is left invertible.

Proof of Theorem 1.6. All that remains to be shown is: T^{α}_{ϕ} is invertible for each $\alpha \in \mathbb{B}^2$ if and only if the distance from ϕ to the invertible elements of \mathfrak{A} is at most one. If T^{α}_{ϕ} is invertible for each $\alpha \in \mathbb{B}^2$, then there exists a $\psi \in \mathfrak{A}$ such that $\|\phi - \psi\| < 1$ by Lemma 3.9. By Lemma 3.7, $T^{\alpha}_{\phi} T^{\alpha}_{\psi}$ is invertible. By Lemma 3.1, T^{α}_{ϕ} is invertible and thus T^{α}_{ψ} is invertible. B Lemma 3.6 ψ is invertible in \mathfrak{A} .

The converse is contained in Lemma 3.7.

References

- Abrahamse, M. B. Toeplitz operators in multiply connected regions, Amer. J. Math. 96 (1974), 261–297. 2
- [2] Abrahamse, M. B. The Pick interpolation theorem for finitely connected domains, Michigan Math. J. 26 (1979), no. 2, 195–203.
- [3] Abrahamse, M. B.; Douglas, R. G. A class of subnormal operators related to multiply-connected domains, Advances in Math. 19 (1976), no. 1, 106–148. 2
- [4] Ahern, P.R.; Sarason, D. The H^p spaces of a class of function algebras, Acta Mathematica, 117 (1967), 123-163.

THE NEIL ALGEBRA

- [5] Ball, Joseph A.; Bolotnikov, Vladimir; ter Horst, Sanne. A constrained Nevanlinna-Pick interpolation problem for matrix-valued functions, Indiana Univ. Math. J. 59 (2010), no. 1, 15–51 2
- [6] Ball, Joseph A.; Guerra Huamn, Moiss D. Convexity analysis and the matrix-valued Schur class over finitely connected planar domains, J. Operator Theory 70 (2013), no. 2, 531–571. 2
- Ball, Joseph A.; Guerra Huamn, Moiss D. Test functions, Schur-Agler classes and transfer-function realizations: the matrix-valued setting, Complex Anal. Oper. Theory 7 (2013), no. 3, 529–575.
- [8] Broschinski, Adam. Eigenvalues of Toeplitz operators on the annulus and Neil algebra, Complex Anal. Oper. Theory 8 (2014), no. 5, 1037–1059. 2, 3
- [9] Conway, John B. A Course in Functional Analysis, Graduate Texts in Mathematics, 96. Springer, New York, 2007. 8
- [10] Davidson, Kenneth R.; Paulsen, Vern I.; Raghupathi, Mrinal; Singh, Dinesh. A constrained Nevanlinna-Pick interpolation problem, Indiana Univ. Math. J. 58 (2009), no. 2, 709–732. 2, 3
- [11] Douglas, Ronald G. Banach algebra techniques in operator theory, Pure and Applied Mathematics, Vol. 49. Academic Press, New York-London, 1972. 2, 7
- [12] Dritschel, Michael A.; Pickering, James. Test functions in constrained interpolation, Trans. Amer. Math. Soc. 364 (2012), no. 11, 5589–5604. 2
- [13] Jury, Michael T.; Knese, Greg; McCullough, Scott. Nevanlinna-Pick interpolation on distinguished varieties in the bidisk, J. Funct. Anal. 262 (2012), no. 9, 3812–3838. 2
- [14] Koosis, Paul. Introduction to Hp spaces. With an appendix on Wolff's proof of the corona theorem, London Mathematical Society Lecture Note Series, 40. Cambridge University Press, Cambridge-New York, 1980. 1
- [15] Martinez-Avendaño, Rubén A.; Rosenthal, Peter. An introduction to operators on the Hardy-Hilbert space, Graduate Texts in Mathematics, 237. Springer, New York, 2007. 6
- [16] Raghupathi, Mrinal. Abrahamse's interpolation theorem and Fuchsian groups, J. Math. Anal. Appl. 355 (2009), no. 1, 258–276 2
- [17] Raghupathi, Mrinal. Nevanlinna-Pick interpolation for C + BH[∞], Integral Equations Operator Theory 63 (2009), no. 1, 103–125. 2
- [18] Sarason, Donald. The H^p spaces of an annulus, Mem. Amer. Math. Soc. No. 56 1965. 2

SRIRAM BALASUBRAMANIAN, DEPARTMENT OF MATHEMATICS, IIT MADRAS, CHENNAI - 600036, INDIA

E-mail address: bsriram@iitm.ac.in

Scott McCullough, Department of Mathematics, University of Florida, Gainesville *E-mail address*: sam@ufl.edu

UDENI WIJESOORIYA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE *E-mail address*: wudeni.pera06@ufl.edu