



Urban air quality management—A review

Sunil Gulia¹, S.M. Shiva Nagendra², Mukesh Khare¹, Isha Khanna¹

¹ Civil Engineering Department, Indian Institute of Technology Delhi, Hauz Khas, New Delhi, India

² Civil Engineering Department, Indian Institute of Technology Madras, Chennai, India

ABSTRACT

Urban air quality management plan (UAQMP) is an effective and efficient tool employed in managing acceptable urban air quality. However, the UAQM practices are specific to a country's needs and requirements. Majority of the developed countries have full-fledged UAQMP with a regulatory management framework. However, developing countries are still working in formulating the effective and efficient UAQMPs to manage their deteriorating urban air environment. The first step in the process of formulation of UAQMP is to identify the air quality control regions based on ambient air quality status and second, initiate a time bound program involving all stakeholders to develop UAQMPs. The successful implementation of UAQMPs depends on the strength of its key components, e.g. goal/objective, monitoring network, emission inventory, air quality modeling, control strategies and public participation. This paper presents a comprehensive review on UAQMPs, being implemented worldwide at different scales e.g., national (macro), city (medium), and local (micro).

Keywords: Urban air quality management plan, urban hot spot, vehicular air pollution, air quality monitoring and modeling, urban air pollution control



Corresponding Author:

Mukesh Khare

☎ : +91-11-26591212

☎ : +91-11-26581117

✉ : kharemukesh@yahoo.co.in

Article History:

Received: 28 April 2014

Revised: 29 September 2014

Accepted: 30 September 2014

doi: 10.5094/APR.2015.033

1. Introduction

Urban air pollution (UAP) is a major concern throughout the world in both, developed and developing countries. Swelling urban population and increased volume of motorized traffic in cities have resulted in severe air pollution affecting the surrounding environment and human health. The World Health Organization (WHO) has estimated that in developing countries, increasing UAP has resulted in more than 2 million deaths per annum along with various cases of respiratory illnesses (WHO, 2005; Cities Alliance, 2007; WHO, 2014). One of the major sources of UAP is the road transport sector. Besides, domestic, commercial and industrial activities also contribute to UAP. It is reported that over 70–80% of air pollution in mega cities in developing nations is attributed to vehicular emissions caused by a large number of older vehicles coupled with poor vehicle maintenance, inadequate road infrastructure and low fuel quality (Auto Fuel Policy, 2002; Molina and Molina, 2002; Badami, 2005; Anjaneyulu et al., 2006; Molina et al., 2007; Singh et al., 2007; Wang et al., 2010). The criteria pollutants responsible for deteriorating urban air quality are oxides of nitrogen (NO_x), sulfur dioxide (SO₂), carbon monoxide (CO), particulate matter (PM) and volatile organic compounds (VOCs). Re-suspension of road dust due to movement of traffic and tyre and brake wear are also some of the significant sources of ambient PM concentrations in urban areas (Amato et al., 2014). Ambient air pollutant concentrations are distributed non-uniformly in urban areas, creating hot spots mostly in central business district, traffic intersections and signalized roadways (Kandlikar, 2007). Besides, topographical and meteorological variations in urban areas lead to complex spatial and temporal variations in pollutant concentrations (Gokhale and Khare, 2007). The spatial scale of UAQMP

varies from macro (national level) to medium (city level) and micro level (site specific). The temporal scale is either long-term or short term, based on the national ambient air quality standard (NAAQS). Further, Table 1 describes the UAQM definitions/concepts (Laxen, 1993; Longhurst et al., 1996; Steinar et al., 1997; Fedra and Haurie, 1999; Beattie et al., 2001; Karatzas, 2002; Gokhale and Khare, 2007; Vlachokostas et al., 2009; Sivertsen and Bartonova, 2012). In addition, Woodfield et al. (2006) have evaluated regional groupings among local authorities to manage urban air quality in London, the West Midlands and former-Avon area of Southwest England. They have reported variations in methods and tools used in developing the local air quality management plan. Further, Longhurst et al. (2009) have discussed the challenges in the source-control approach of air quality management and recommended the development of an integrated, risk management effect-based process of urban air quality management. They have also reported the importance of periodically evaluating the dynamic nature of management challenges by reviewing components of UAQM protocol. In the recent past, Figueiredo et al. (2013) have carried out an assessment of source contribution to urban air quality in the city of Estarreja in Portugal and observed that O₃ and PM₁₀ concentrations exceeded urban air quality standards. As a result, the authors have suggested strategies to reduce O₃ and PM₁₀ concentrations from motorized road transport and industrial activities. A majority of the existing UAQMPs have been developed both at macro (national) or medium (city) levels and for long-term duration considering the average ambient pollutant concentrations (Steinar et al., 1997; NRC, 2004; CPCB, 2006; NILU, 2007; DEAT, 2008). Gokhale and Khare (2007) have introduced the concept of an e-UAQMP framework to forecast the air pollution episodes at a selected urban air quality control region and further suggested

mitigation plans. This paper is an attempt to present a comprehensive review on UAQMPs that assist regulatory authorities in maintaining the acceptable urban air quality.

2. Urban Air Pollution in Mega Cities

In developed countries, trends of urbanization and the associated growth of cities have started to reverse due to severe levels of congestion (Mayer, 1999). However, in developing countries, city's growth tends to be from periphery to core. The ambient air pollution levels at urban hot spots in 20 European cities have exceeded the urban background concentrations due to increase in traffic volume (Moussiopoulos et al., 2005). In the UK, motorized road transport has been categorized as one of the largest single pollution sources in 92% of declared air quality management areas (AQMPs) which accounts for 33% emissions of NO_x and 21% of PM_{10} , and so, frequently violating the national ambient air quality standards/objectives (Faulkner and Russell, 2010). However, in the recent past, it has been observed that in some mega cities of developed countries, urban air quality is showing signs of improvement on account of efficient implementation of UAQMP (NSW Government, 2010; EEA, 2011a; Parrish et al., 2011; EEA, 2013a). In the European Environment Agency (EEA) countries, the emission reduction from vehicular exhausts from 1990 to 2009 has been reported to be around 54% for SO_2 , 27% for NO_x , 16% for PM_{10} and 21% for $\text{PM}_{2.5}$. In spite of all these efforts in place, it has been observed that 18% to 49% of the population in these countries is still exposed to PM_{10} concentrations exceeding the ambient standards (EEA, 2011a). In North American megacities like, Los Angeles, New York, and Mexico City, the ambient air quality concentrations for some criteria pollutants have shown declining trends during the last five decades, particularly the O_3 concentrations. In the year 2010, 8-hour average ambient O_3 concentrations have been observed to be 17% lower compared to the concentrations in the year 1990. Similarly, 24-hour average PM_{10} concentrations have been found to be 38% lower in 2010 when compared with levels in 1990; and so with NO_2 and SO_2 concentrations which have been reported to be 45% and 17% lower in 2010 when compared with 1990 concentration levels, respectively. However, at some designated non-attainment areas NAAQS is still violated (Parrish et al., 2011; U.S. EPA, 2012). In New South Wales (NSW) in Australia, road transport is the single largest source of NO_x emissions that contributes more than 71% of total emissions. One-hourly average NO_2 concentrations have shown a declining trend from 1980 to 2009. However, the annual average

concentration trend of $\text{PM}_{2.5}$ has remained more or less constant from 1997 to 2009. The reduction in ambient NO_2 concentrations during this period may be due to the implementation of UAQMP maintaining cleaner fuel standards (NSW Government, 2010). The cited examples clearly show definite benefits of the UAQMPs.

In the Asian subcontinent, some developed countries, e.g. Singapore, Japan and Hong Kong, are facing street-level air pollution problems due to an increase in the number of motorized transport (ADB, 2006a; CEC, 2011; Edesess, 2011). In developing countries, mega cities are facing acute problems due to an increase in the ambient PM and NO_2 concentrations as a result of rapid urbanization. In Shanghai, New Delhi, Mumbai, Guangzhou, Chongqing, Calcutta, Beijing, and Bangkok the ambient PM and NO_2 concentrations frequently violate WHO guidelines (Baldasano et al., 2003). Poor fuel quality, high traffic density, large proportion of old vehicles, poor road infrastructure and inadequate inspection and maintenance (I/M) programs, are some of the major causes of deteriorating urban air quality (Gurjar et al., 2004; Badami, 2005). Chan and Yao (2008) have reported that ambient concentrations of PM_{10} and SO_2 in the Chinese cities of Shanghai and Pearl River Delta are four to six times higher than concentrations observed in any of the cities in developed countries. In Beijing, the annual average NO_2 concentrations remain constant at a level of $70 \mu\text{g}/\text{m}^3 \pm 10\%$. However, 90% of the time, PM concentrations exceed the NAAQS and WHO-AQG (Zheng et al., 2005; Duan et al., 2006). One of the studies carried out recently has found that the annual average PM_{10} concentrations in Asian cities are four times higher than WHO-AQG of $20 \mu\text{g}/\text{m}^3$ (Atash, 2007; CAI-Asia, 2010a; UN-Habitat, 2010). In the Indian metropolitan cities (Delhi, Mumbai, Kolkata and Chennai), ambient PM concentrations frequently violate NAAQS as well as WHO guidelines (Gupta and Kumar, 2006; Singh et al., 2007; CPCB, 2010a; Gupta et al., 2010). Mohan and Kandya (2007) have analyzed nine years of data at seven different locations in the city of Delhi and reported that at one of the locations (ITO intersection), the air quality has been found to be "worst". In South Africa, the air quality act, 2004 contains specific provisions to deal urban hot spot by declaring it as the "priority area" (DEAT, 2008). Gurjar et al. (2008) have developed a multi-pollutant index for 18 megacities of the world, out of which, five are classified as "fair" and thirteen as "poor". Further, Ramachandra and Shwetmala (2009) have reported that India's transport sector emits 258.10 Tg of CO_2 , of which 94.5% is contributed by motorized road transport. The Central Pollution Control Board (CPCB) has reported that vehicular contribution to the total urban air pollution

Table 1. The UAQM definition/concept

References	Air quality management definition/concept
Laxen (1993)	AQMP is a program to ensure the control of emissions to protect public health and welfare.
Longhurst et al. (1996)	Concept of the local air quality management plan (LAQMP) emphasizes the distribution of power at community level and strong communication and cooperation between actors of air quality management.
Steinar et al. (1997)	AQMP is a system for design and implementation of monitoring, management and policies within an urban area.
Fedra and Haurie (1999)	A system that can establish a robust and integrated environmental management life cycle for urban areas of interest.
Beattie et al. (2001)	The UAQM emphasizes on local action to deal urban hot spots of air pollution (episode).
Karatzas (2002)	A software tools that combine air quality models with other software modules like geographical information systems, databases, expert systems and statistical analysis tools. It helps to interpret the complex interactions between various atmospheric, emission, land use and topographic parameters.
Woodfield et al. (2003)	The UAQM identifies specific pollution hot spots based on advanced and complex air-quality dispersion modeling and monitoring techniques.
Gokhale and Khare (2007)	Episodic urban air quality management plan (e-UAQMP)—it uses hybrid model to forecast extreme levels of pollutants during peak traffic hours.
Vlachokostas et al. (2009)	AQMP is an integrated assessment methodological scheme for the evaluation of air pollution control measures that are put forward in order to reduce air pollution levels in urban areas. This approach brings together air quality modeling and mathematical programming techniques, provides a decision support system for the determination of optimal bundles of air pollution control options according to the particular features and needs of the areas examined.
Sivertsen and Bartonova (2012)	The UAQM is a set of action that helps in attaining air quality goals in a specified geographical area. It requires actions by government, business, industry, NGO's and the population.

in Delhi and Mumbai is about 76–90% for CO, 66–74% for NO_x, 5–12% for SO₂ and 3–12% for PM (CPCB, 2010a). Recently, studies carried out by Yale University, USA, and WHO, have ranked Delhi as the “worst” polluted city based on environment performance index (TOI, 2014; The Hindu, 2014). Therefore, necessary mitigation measures need to be implemented through effective and efficient implementation of UAQMP to maintain an acceptable urban air quality.

3. Urban Air Quality Management

The UAQM practices are country specific and based on the priorities as agreed for a specific air quality control region to maintain an acceptable ambient air quality. They are implemented and enforced through legislative laws (Elsom, 1996; Longhurst et al., 1996). Table 2 describes and compares various legislative laws and regulations dealing with UAQM which have been implemented in selected developed and developing countries of the world. The key components of UAQMP are air quality objectives, monitoring, emission inventory, prediction and forecasting tools, control strategies and public participation. Further, each component plays a significant role in improving the efficiency of the UAQMP, thus reducing pollutant concentrations. Moreover, the effective and efficient implementation of UAQMP in developing countries still remains a challenging task for air quality managers due to lack of government commitment and stakeholder participation, weaknesses in policies, standards and regulations, lack of real-time air quality data and emission inventories (KEI, 2002; ADB, 2006b; Naiker et al., 2012). In one of the studies carried out in Indonesia, Santosa et al. (2008) have reported that urban air pollution is perhaps the most severe environmental problem due to rapid growth in industrial and transportation sectors. The management practices to improve urban air quality are very limited and the portion of the budget allocated for urban air quality management is also insufficient. In the recent past, Kura et al. (2013) have analyzed urban air pollution problems in China, India and Brazil at a macro urban scale and proposed a system based methodology to develop the UAQMP that takes into account (i) identification of critical pollutants and their sources, (ii) setting up of the air quality monitoring network, (iii) emission inventory, (iv) source prioritization, (v) control strategies, and (vi) development of decision support system. Key components of the existing UAQMP have been critically reviewed and discussed in the following sections.

3.1. Air quality objectives/standards

Air quality objectives/standards are principal components of any UAQMP that act as a prerequisite to clear policies and goals. The formulation of the objectives/standards requires comprehensive discussions and reviews with stakeholders at various levels of hierarchy e.g. national, regional and local (Longhurst et al., 1996). In USA, air quality standards are categorized for ambient and hazardous pollutants. The ambient standards are described under NAAQS consisting of criteria pollutants such as O₃, CO, SO₂, NO₂, and PM (U.S. EPA, 2011a). In the UK, the ambient air quality standards are based on the carrying capacity of the environment. However, the UAQM policies specify a target timeline within which the controlling authorities have to achieve the objectives/standards. The EU's air quality directive contains limit and target values (DEFRA, 2011a).

In the developing world, most of the countries have evolved their own air quality standards. However, some countries follow the WHO–AQG and specify them as their standards (WHO, 2005). The WHO–AQG for PM is stringent than any other ambient air quality standards specified in USA, UK and EU countries. In South Africa, the air quality objectives and standards are formulated involving stakeholders and community goals and priorities (DEAT, 2008). In India and China, the NAAQS has been formulated based on land use type i.e. sensitive, residential/commercial and indus-

trial (MoEF, 2009; MEP, 2013). NAAQS in India has been set up for 13 pollutants (MoEF, 2009). Table 3 describes air quality standards/objectives followed in different countries of the world including the WHO–AQG.

3.2. Air quality monitoring

Air quality monitoring provides information regarding the status of present air quality. It helps in evaluating the existing policies and their effective implementation. One of the important components of any air quality monitoring program is planning, design and establishment of monitoring network based on the air quality objectives (Sivertsen, 2008). Graves et al. (1981) have designed a monitoring network in Fulton County, Georgia, in which only non-reactive pollutants are monitored in ambient air. Air quality monitoring network in Greater London area has been designed specifically with an objective to carry out spatial correlation analysis using the data obtained from the monitoring stations (Handscombe and Elsom, 1982). Mofarrah and Husain (2010) have used multiple-criteria approach in designing an optimal air quality-monitoring network in Riyadh City, Saudi Arabia. Table 4 describes various air quality-monitoring networks being used in different countries.

In USA, the monitoring network is designed and operated following common guidelines of Office of Air Quality Planning and Standards. Monitoring network data are analyzed to forecast air quality index (AQI) in over 300 cities (AIRNow, 2011). In UK, over 300 air quality-monitoring stations are operated (DEFRA, 2013). Real time hourly average air quality information is available through electronic media and web platforms. The EU countries administer their monitoring network through European Topic Centre on Air Pollution and Climate Change (EEA, 2014).

In most of the developing countries, the air quality-monitoring network has been designed primarily to ensure effective regulatory compliance. In South Africa, 94 air quality-monitoring stations are operated throughout the country ensuring regulatory compliance (DEA, 2011). In China, the CNEMC is responsible for operation and maintenance of air quality monitoring stations in 113 cities with more than 2 000 stations (Shasha, 2010). In Hong Kong, 14 online continuous monitoring stations are in operation and maintained by Environment Protection Department (EPD, 2011). However, the Pearl River Delta regional air quality monitoring network serves as a role model for cooperation between the two governments and among various local authorities to address the air pollution problems in a more effective and holistic manner (Zhong et al., 2013a). In India, national ambient air quality monitoring (NAAQM) network is having 342 monitoring stations in 127 cities/towns in 26 states and 4 union territories. Additionally, individual state pollution control boards are operating their own monitoring stations (CPCB, 2014). In the recent past, the Ministry of Earth Sciences, Government of India has started monitoring of criteria air pollutants, specifically designed for Delhi City and operated by the Indian Institute of Tropical Meteorology (IITM), Pune (IITM, 2014).

3.3. Source apportionment

Source apportionment (SA) studies are conducted to identify and quantify the impact of different sources of air pollutants at receptor sites. An efficient and effective UAQMP needs input on categories of sources that may contribute to ambient air pollution followed by their quantification. Based on this information, effective UAQM strategies can be formulated and implemented. The SA studies can be performed using methods that rely on an analysis of morphological and chemical composition of pollutants. The latter, being the quantitative technique, is preferred over the former one for use in large-scale studies. Chemical characterization, thus, is an important step in the SA studies that focuses on obtaining chemical constituents of the PM, which depend on

Table 2. Urban air quality management system/process

Country	Air Quality Management Process	Scale	Designated Area	References
USA	The management practices were introduced in the Clean Air Act (CAA), 1963 and Air Quality Act (AQQA), 1967. The CAA, 1963 recommends a set of responsibilities and relationships among federal, state, tribal, and local agencies regarding air quality management. The USEPA and states have designated the non-attainment areas (NAAs) that experience <i>unhealthy</i> levels due to exceedences of <i>criteria</i> pollutants. The local governments have developed rules/guidelines (State Implementation Plan, SIP) to reduce the pollution level in order to comply with (or attain) the specified standard.	Local and district level	NAAs	NRC (2004); Green and Cummins (2007)
Europe	The air quality management practices were introduced in 1980 with directive 80/779/EEC. In UK, the LAQM technical guidelines support local authorities in carrying out their duties under the Environment Act 1995 and the Environment (Northern Ireland) Order 2002 which further revised and improved in 2000, 2003 and 2009 with participation of stakeholders.	Area specific e.g. urban <i>hot spot</i>	AQMA	Longhurst et al. (1996); DEFRA (1998); DEFRA (2009a); DEFRA (2009b)
Australia	Urban air quality is managed through the national plan instead of region-specific plan of Environment Protection and Biodiversity Conservation Act, 1999. City councils have their own UAQMPs with more emphasis on emission reduction from transport sector.	National/city scales	AQMA	EPA (2007); Commonwealth of Australia (2011); AQCC (2012)
British Columbia	The UAQMP is implemented through Clean Air Provisions of Environmental Management Act, 2004. The air pollution management is applied successfully as airshed action strategies.	Local scale (airshed)		Williams and Bhattacharyya (2004); B.C. Government (2008)
Mexico	UAQM practices are implemented first time in 1999 to reduce the criteria pollutant concentrations in urban areas. The ministry of environment and natural resources is the regulatory body to implement the UAQM strategies throughout Mexico.	City scale	NAAs	Molina and Molina (2002)
South Africa	The National Environmental Management–Air Quality Act, 2004 introduced the UAQMP. This act implemented air quality standards in South Africa. The primary aim is decentralization of the implementing agencies.	City scale	Priorities area	DEAT (2008); Naiker et al. (2012)
China	The UAQM starts with 2 nd revision in Air Pollution Prevention and Control Act in 2000. Recently, the state council has introduced action plans to control the urban air pollution at city as well as local scales.	City/local scale		Chan and Yao (2008); Wang and Hao (2012); CAAC (2013); MEP (2013)
India	The UAQM practices initiated after implementation of the Air (Prevention and Control of Pollution) Act, 1981. At present, no UAQMP exists to manage urban air pollution in Indian cities. The CPCB has issued guidelines to prepare action plans for sixteen polluted cities.	City scale	NAAs	CPCB (2006)

Table 3. Comparative analysis of NAAQS for criteria pollutants

Pollutants	Developed Countries			Developing Countries					International			
	UK/EU 2005 ^a	US ^b	Australia ^c	British Columbia ^d	South Africa ^e	Mexico ^f	China I ^g	China II ^g	China III ^g	India I ^h	India II ^h	WHO ⁱ
1 hr	350, 24/year ^j		200	900	350, 88/year	SO ₂ (µg/m ³)	150	500	700			
24 hr	125, 3/year	365, 1/year	80	260	120, 4/year	0.13 ppm 1/year	50	150	250	80	80	125
Annual	20		20	50	50 0.03 ppm	NO ₂ (µg/m ³)	20	50	100	50	20	50
1 hr	200, 18/year, (EU 2010)		120	400	200, 88/year	0.21ppm, 1/year	120	120	240			200
24 hr				200	40		80	80	120	80	80	
Annual	40 (EU, 2010)	100	30	60	40		40	40	80	40	30	40
24 hr	50, 35/year		50	50	120, 4/year	PM ₁₀ (µg/m ³)	50	150	250	100	100	50
Annual	40	50			50		40	100	150	60	60	20
24 hr		35, 98%	25	25	65		50	150	250	60	60	25
Annual	25	15	8	8	15		40	100	150	40	40	10
1 hr		40 000, 1/year		28 000	30 000, 88/year	CO (µg/m ³)	10 000	10 000	10 000	4 000	4 000	30 000
8 hr		10 000, 1/year	9 000	11 000	10 000, 11/year	11 ppm 1/year	4 000	4 000	2 000	2 000	2 000	10 000
24 hr							4 000	4 000	4 000			

^a Source: http://uk-air.defra.gov.uk/documents/National_air_quality_objectives.pdf

^b Source: <http://www.epa.gov/air/criteria.html>

^c Source: <http://www.environment.gov.au/topics/environment-protection/air-quality/air-quality-standards#air>

^d <http://www.bcairquality.ca/reports/pdfs/aqatable.pdf>

^e DEA (2009)

^f http://transportpolicy.net/index.php?title=Mexico_Air_Quality_Standards (PM₁₀ and PM_{2.5} value should be met at 98% of the time in a year)

^g China: Zone I: Residential areas; Zone II: Commercial areas; Zone III: Industrial areas (http://www.vecc-sepa.org.cn/eng/news/news_detail.jsp?newsid=e00397)

^h India I -Industrial, Residential, Rural and other area, India II -Ecological sensitive area, 98 % of the time in year

ⁱ WHO-AQG (WHO, 2005)

^j 24 per year means 24 exceedances per year are permitted

Table 4. Air quality monitoring networks

Countries	Air Quality Monitoring Network	Automatic/Manual	Pollutants Monitored	Available Online
US	(i) State and local air monitoring stations (SLAMS), (ii) National air monitoring stations (NAMS) (iii) Special purpose monitor station (SPMS) for very specific or short-term monitoring goals and (iv) Photochemical assessment monitoring station (PAMS). Categorized based on land use type: Urban (U), Suburban (S), Rural (R), Traffic (T), Industrial (I) and Background (B). Automatic Rural and Urban Network (ARUN) is the largest monitoring network.	Both	CO, Pb, NO ₂ , O ₃ , PM, SO ₂	www.airnow.gov
UK		Both	NO _x , SO ₂ , O ₃ , CO and PM (PM ₁₀ and PM _{2.5})	www.uk-air.defra.gov.uk/network
EU	Categorized based on compliance, exposure assessment, on-line monitoring (episode), and operational monitoring (adjacent to specific source).	Both	NO _x , NO ₂ , SO ₂ , O ₃ , CO, PM (PM ₁₀ and PM _{2.5}), Benzene	www.eea.europa.eu/data-and-maps/data/airbase
Australia	Urban and Regional based on population >25000. Monitoring is not done if previous measurements or screening studies have shown that specific pollutant levels would be consistently below the NAAQS.	Both	CO, NO ₂ , O ₃ , SO ₂ , PM ₁₀ and PM _{2.5}	www.environment.nsw.gov.au/aqms
British Columbia	Continuous (hourly), non-continuous (24 hourly) and mobile (episodic).	Both	CO, NO ₂ , O ₃ , PM _{2.5} , PM ₁₀ , SO ₂	www.envistaweb.env.gov.bc.ca
Mexico	Manual monitoring stations started in 1972. Automatic air-quality monitoring network established in late 1980s. Air quality data are shared through National Information System.	Both	O ₃ , NO ₂ , SO ₂ , PM ₁₀ , PM _{2.5} , CO	http://sinaica2.inecc.gob.mx/magic/rmart
South Africa	The South Africa Air Quality Information System (SAAQIS).	Manual	CO, NO ₂ , PM ₁₀ , SO ₂	www.saaqis.org.za
China	The Chinese National Environmental Monitoring Center (CNEMC).	Both	CO, SO ₂ , O ₃ , NO ₂ and PM ₁₀ and PM _{2.5}	www.aqicn.org
India	National Ambient Air Quality Programme (NAAQM).	Both	SO ₂ , NO _x , SPM, CO and PM ₁₀	http://164.100.43.188/cpcbnew

sources and their emission rates. To apportion the sources, receptor models are used, which quantify pollutant concentrations based on the measured ambient air pollutant data. SA studies using receptor models date back to late 1960's, with the first study reported in literature by Blifford and Meeker (1967). They have examined particle composition data of SPM collected by the National Air Sampling Network (NASN) during 1957–61 in 30 U.S. cities using factor analysis and eigenvector method for 13 elements. Very few studies have been conducted nation-wide (Thurston et al., 1984; CPCB, 2010a; Thurston et al., 2011); majority of the studies have been conducted only on regional and local scales (Ito et al., 2004; Qin et al., 2006; Tao et al., 2013; Li et al., 2014). However, the focus of these studies has been shifting from coarser particles (SPM, PM₁₀) to finer particles (PM_{2.5}, PM₁) that significantly affect health (Schauer and Cass, 2000; Mazzera et al., 2001; Chakrobarty and Gupta, 2010; Khare and Baruah, 2010; Pandolfi et al., 2011). One of the important parameters in SA studies is the selection of marker species or source profiles. Metals have been used invariably as marker species for identification of sources—for instance, Al, Si, Ti, Ca for crustal/soil sources; Ni, V for residual/fuel oil combustion; Zn, Cr for refuse burning/incineration; and Zn, Cr, Pb, Cu for vehicular emissions (Watson et al., 2001; Chow et al., 2011; Patil et al., 2013). The ratio between two metal concentrations represents the characteristic of a particular source which is an important input for receptor models. Besides, an efficient and effective UAQMP requires specific information on emission sources which may be categorized based on the type of fuel used. In order to identify such sources, organic molecular markers are used as tracers along with metals, ions and EC/OC (Schauer et al., 1996; Perrone et al., 2012; Li et al., 2014). These markers have a high degree of source specificity; for instance, levoglucosan from cellulose has been used as a specific marker compound for wood burning; hopane and stearanes for mobile source emissions; and polycyclic aromatic hydrocarbon (PAH) profiles for distinguishing between gasoline and diesel vehicle emissions (Simoneit et al., 1999; Sharma et al., 2003; Chowdhury et al., 2007; Lin et al., 2010; Masih et al., 2010; Herlekar et al., 2012). Hasheminassab et al. (2013) have used molecular marker-based chemical mass balance model (MM-CMB), which possesses higher source identification efficiency when compared to CMB, which is one of the most widely used receptor models (Watson et al., 2002; Watson et al., 2008). Other receptor modeling techniques used in SA studies include enrichment factor analysis, times series analysis, multivariate factor analysis (principal component analysis and positive matrix factorization), UNMIX, species series analysis and multi-linear engine (ME) analysis (Cooper and Watson, 1980; Henry et al., 1984; Hopke, 1991; Ramadan et al., 2003; Almeida et al., 2006; Hwang et al., 2008; Viana et al., 2008; Gietl and Klemm, 2009; Begum et al., 2010; Kertesz et al., 2010; Oliveira et al., 2010; Stone et al., 2010; Begum et al., 2011; Cheung et al., 2011; Pakbin et al., 2011; Tecer et al., 2012; Daher et al., 2013; Bove et al., 2014; Kuo et al., 2014). However, few SA studies have also been carried out using dispersion modeling approach (Colville et al., 2003; Laupsa et al., 2009). Recent studies carried out in some of the urban areas in USA have concluded that metal/steel industries, motor exhaust and crustal emissions are major contributors of PM_{2.5} concentrations in ambient air (Coutant et al., 2003; Thurston et al., 2011; Green et al., 2013; Hasheminassab et al., 2013; Sturtz et al., 2014). Studies conducted in European urban cities, motorized transport, crustal sources and mixed industrial/fuel-oil combustion are major sources of PM_{2.5} and PM₁₀ (Viana et al., 2008; Yin et al., 2010; Belis et al., 2013).

Since the last decade, many developing countries have included SA as an important component of UAQMP. In China, several studies have been carried out focusing identification, categorization and quantification of PM sources in ambient urban air and their correlation with human health (Song et al., 2006; Xie et al., 2008; Gu et al., 2010; Kong et al., 2010; Breitner et al., 2011; Gu et al., 2011; Leitte et al., 2011; Zhu et al., 2011; Liu et al., 2013; Wang et al., 2013). In India, a number of SA studies have been

carried out apportioning SPM, PM₁₀ and PM_{2.5}, using receptor modeling techniques (Khillare et al., 2004; Chowdhury et al., 2007; Gupta et al., 2007; Sharma et al., 2007; Chelani et al., 2008; Kothai et al., 2008; Srivastava and Jain, 2008; Srivastava et al., 2008; CPCB, 2010a; Kar et al., 2010; Shridhar et al., 2010; Gummeneni et al., 2011; Srimuruganandam and Nagendra, 2011; Pant and Harrison, 2012; Pant and Harrison, 2013; Tiwari et al., 2013; Sharma et al., 2014). One such similar study has recently been completed by CPCB, a regulatory body in India, covering six mega cities and presenting an integrated approach of measurements, chemical speciation and receptor modeling for assessing efficacy of UAQMP strategies in mitigating PM₁₀ ambient levels (CPCB, 2010a). Further, SA studies also provide a platform for the development of comprehensive and accurate emission inventories.

3.4. Emission inventory

A comprehensive emission inventory is the first step to develop an emission control strategy for selected pollutants. Compilation of an accurate emission inventory is an integral part of any UAQMP (Moussiopoulos, 2003; Miller et al., 2006). The product of the emission factor and traffic volume provides emission rate of motorized traffic (Esteves-Booth et al., 2002). Emission factors are normally expressed in grams of a pollutant per unit of distance traveled. The emission factor depends on the type of vehicle and fuel used, type of engine, I/M program, vintage of vehicles, driving cycle and average speed (Auto Fuel Policy, 2002). Emission inventory models are available, which allow users to vary the fleet structure, technology proportions, vehicle activity and proportions of driving conditions to estimate emissions and total fuel consumption (Boulter et al., 2007). U.S. EPA (2011b) describes various emission models including MOVES, MOBILE-6 and NMIM. Recently, Kota et al. (2014) have evaluated emission inventory of CO and NO_x from on-road vehicles using MOBILE 6.2 and MOVES in Southeast Texas, USA. They have found that MOBILE 6.2 estimations are more accurate when compared to MOVES. In UK, DEFRA (2011b) provides and maintains national atmospheric emission inventory. A majority of the EU countries use COPERT to estimate the emission load from motorized vehicles (Ekstrom et al., 2004). However, in Germany and Switzerland, "Handbook of Emission Factors" is also used for estimating the emission load from motorized sources (Keller and Zbinden, 2004). In Mexico, Wolf et al. (2009) have developed National Emissions Inventory for criteria pollutants and reported that mobile sources are responsible for 16% of PM₁₀, 52% of PM_{2.5}, 99% of CO, 82% of NO_x, 31% of NMVOC (non-methane volatile organic compounds) and 22% of ammonia (NH₃) emissions.

In China, MOBILE5B-China is used for motor vehicle emission inventories that are based on US EPA's MOBILE5b and PART5 (Hao et al., 2000; Hao and Wang, 2005; Zhou et al., 2010). Huo et al. (2009) have proposed a new link-based, bottom-up vehicle emission inventory method for Beijing City using the available on-road emission measurement data and activity survey data. Results have stated that total emissions from light-duty vehicles in Beijing urban area in 2004 have been 1 141 Mg of CO per day, 48 Mg of HC per day, and 32 Mg of NO_x per day. In South Africa, emission rates are estimated by simple multiplication of emission factor and traffic volume (DEAT, 2008). In India, the Indian Institute of Petroleum has estimated emission factors for two wheelers and cars based on standard Indian driving cycle (Pundir and Das, 1985). Further, the Automotive Research Association of India, in collaboration with CPCB, New Delhi has developed emission factors for all motorized vehicle types based on standard driving cycle and average vehicle speed (ARAI, 2007). In another study, Nagpure and Gurjar (2012) have developed a vehicular air pollution inventory (VAPI) model for Indian traffic conditions. In the recent past, Guttikunda and Calori (2013) have developed a GIS based emission inventory of air pollutants in Delhi City.

3.5. Air quality modeling

Air quality modeling plays an important role in formulating air pollution control and management strategies by providing guidelines for efficient air quality planning. Its main objective is to predict ambient air pollutant concentrations of one and more species in space and time as related to independent variables such as emission and meteorological parameters.

In USA, EPA recommended dispersion model, e.g. AERMOD (steady state dispersion model and an advanced version of ISC3) is used for regulatory purposes (Cimorelli et al., 2005; U.S. EPA, 2005). Another EPA recommended model, CALPUFF, is used for long-range transport (source-receptor distances of 50 km to several hundred kms.) of emissions from point, volume, area and line sources (Scire et al., 2000). CALINE 4 is used for highway sources in both urban and rural areas (Benson, 1984). In UK, ADMS-urban is used to assess the ambient air quality (Carruthers et al., 2000). In addition, CMAQ (Byun and Schere, 2006) is used to assess the ambient air quality (Williams et al., 2011). In developing countries, the availability of precise input data is one of the challenges (SEI, 2004). In spite of this, AirQuis modeling system has been developed in South Africa, which is being used for UAQMP in the city of eThekweni (NILU, 2007). Besides, in the designated airshed priority areas, CALPUFF is used to forecast the ambient air quality as a part of UAQMP (DEAT, 2008). At a local scale, Zunckel (2009) has developed an air pollution information network, which comprises guidelines/recommendations on how to carry out efficient air quality modeling. However, in Mexico City, coupled WRF-Chem (a photochemical model) is mainly used to forecast the pollutant concentrations (Tie et al., 2007; Ying et al., 2009; Li et al., 2011). In India, Gaussian based dispersion air quality models are used for regulatory compliances of ambient air quality (CPCB, 1998; CPCB, 2010a). In addition, a number of air quality models/codes have been validated/developed to predict ambient air quality in selected air quality control regions (Luhar and Patil, 1989; Khare and Sharma, 1999; Gokhale and Khare, 2005; Kesarkar et al., 2007; Khare et al., 2012; Nagendra et al., 2012; Sharma et al., 2013a; Sharma et al., 2013b). In addition, Elbir (2004), Elbir et al. (2010), Schipa et al. (2011), Gulliver and Briggs (2011) have used geographical information system (GIS) coupled with air quality models to forecast the pollutant concentrations in ambient urban environments. GIS based forecasting can estimate ambient air pollution levels at high temporal and spatial resolution, generate mapping and scenario results, which can further be compared with air quality standards. This can be integrated with socio-economic data at the receptor site, which may further assist in air pollution related exposure and health assessment and making the UAQMP more productive.

3.6. Air pollution exposure and health assessment

The ultimate goal of any urban air quality management plan is to ensure that the impact of air pollution on human health remains minimal. Past studies have reported that increasing PM concentrations deteriorate human health by affecting respiratory and cardiovascular systems (Dockery et al., 1993; Preutthipan et al., 2004; WHO, 2005; Pope and Dockery, 2006; Colais et al., 2012; Mustafic et al., 2012; WHO, 2014). Therefore, air pollution exposure and health assessment becomes one of the significant components of UAQMP. Pope and Dockery (2006) have reviewed health effects of fine PM on humans and concluded that these effects are functions of concentrations as well as duration of the exposure. They have also reported that long-term exposure studies are more accurate in analyzing the cumulative effects than short-term exposure studies. In the USA, a few studies conducted in the past, have focused on the health impact of air pollution exposure in urban areas (Dockery et al., 1993; U.S. EPA, 2002; Lipfert et al., 2006; Dockery, 2009). In the UK, air pollution exposure assessment is one of the essential components of LAQMP (DEFRA, 2009a). In the European countries, studies have reported

that high air pollution exposure level deteriorates the health of human beings (Gehring et al., 2013; Pascal et al., 2013; Raaschou-Nielsen et al., 2013). In one of the studies done in Australia, it has been observed that motor vehicle exhaust is responsible for causing cardiovascular and respiratory diseases (BTRE, 2005). Simpson et al. (2005) have observed 1% increase in daily total number of deaths per 10 mg/m³ increase in PM_{2.5} concentration. However, in developing countries, very few air pollution exposure and health assessment studies have been conducted so far (Han and Naeher, 2006). Steinar et al. (1997) have suggested incorporation of air pollution exposure and health assessment as an important component of UAQMP for developing countries in the Asian continent. Past studies indicate evidences of high risk of respiratory and cardiovascular diseases due to increased air pollution exposure levels (Terblanche et al., 1992; HEI, 2004; Wichmann and Voyi, 2005; Agarwal et al., 2006; Han and Naeher, 2006; Norman et al., 2007; Gurjar et al., 2010; HEI, 2010). Further, in South Africa, air pollution exposure and health assessment is not included in UAQMP, though many researchers have individually carried out exposure and health impact assessment studies (Wichmann and Voyi, 2005; Norman et al., 2007; DEAT, 2008). One of the studies carried out at Vaal triangle area (a notified priority area) have reported that out of the total surveyed population, 66% suffered from upper respiratory tract illnesses and 29% from lower respiratory tract illnesses (Terblanche et al., 1992). In another study conducted in the urban areas of South Africa, it has been estimated that ambient air pollution levels cause 3.7% of the national mortality from cardiopulmonary disease and 5.1% of mortality attributable to cancers of trachea, bronchus and lung in adults aged 30 years and older; and 1.1% of mortality from acute respiratory infections in children under 5 years of age (Norman et al., 2007). Huang et al. (2009) have analyzed the relationship between air pollution level, visibility and daily mortality rate in Shanghai, China. They have found that an increase in PM concentrations is significantly correlated with reducing visibility and increasing daily mortality rate. They have also suggested that visibility can act as surrogate of air quality in exposure and health assessment. In India, Chhabra et al. (2001) have described the role of ambient air pollution in chronic respiratory morbidity in residents of Delhi City. They have found positive relationship between increasing PM concentrations with chronic respiratory morbidity (i.e., chronic cough, phlegm, breathlessness, bronchitis and bronchial asthma). In another study, HEI (2010) have reported that an increase in PM₁₀ concentrations has increased risk of mortality in Chennai and Delhi cities of India; i.e., 0.4% and 0.15% increase per 10 µg/m³, respectively. It is equivalent to mortality risk associated with PM₁₀ exposure as observed in the first four studies conducted under PAPA (Public Health and Air Pollution in Asia) as well as in multi-city studies conducted in South Korea, Japan, Europe, and North America. Balakrishnan et al. (2011) have proposed a methodology using time-series analysis to quantify the effect of short-term exposure of PM₁₀ on mortality in Chennai City, India. However, it can be concluded that developing countries need to focus more on source-specific exposure assessment studies especially for PM, because different sources of PM have different physio-chemical compositions and thus different biological potentials (Laden et al., 2000; Janssen et al., 2002).

3.7. The UAQM strategies

The UAQM strategies direct their efforts primarily to maintain acceptable ambient air quality in urban corridors by implementing different control measures, such as strict emission norms, improvement in fuel quality, efficient engine technology, I/M program, traffic fleet management within the specified air quality control regions and effective road and transport planning. The UAQM strategies are generally evaluated and selected on the basis of change in pollutant concentrations using air quality dispersion models. However, Sonawane et al. (2012) have reported that exposure and health impacts are more efficient criteria in selecting UAQM strategies. The authors have described that prior to imple-

mentation of UAQM strategies, it should be evaluated in terms of socio-economic and health consequences. The socio-economic evaluation studies need to take into account not only the cost of implementation of UAQM strategies, but also the consequent benefits arising from the actions such as health cost reduction. Steinar et al. (1997) have further stated that before implementation of UAQM strategies, cost-benefit analysis (CBA) or cost effectiveness analysis (CEA) must be carried out. Lieman et al. (2007) and Scorgie (2012) have described qualitative assessment and prioritization of UAQM strategies based on environmental benefits; technical viability; social acceptability and desirability; economic feasibility; and timeframe for implementation and environmental benefits realization. In the UK, after declaration of AQMA, all possible UAQM strategies are evaluated in terms of emission reduction, air quality improvement and cost effectiveness (DEFRA, 2009a; DEFRA, 2009b).

In USA, vehicular emissions are controlled through voluntary mobile source emission reduction programs under SIP (Wilson, 1997). The U.S. Department of Energy (2011) has launched a “Clean Cities Five Year Strategic Plan” aiming at reducing the petroleum-dependence in on-road transportation. In the UK and EU countries, the UAQM strategies are effectively implemented and regularly monitored at specific AQMAs (Corporation of London, 2003; Newcastle City Council, 2006; DEFRA, 2011c; Duffield, 2011; EEA, 2011b; EEA, 2013b; EEA 2013c). However, in developing countries, these are discreetly applied in various forms, e.g. mandatory use of clean fuel (CNG/LNG), use of catalytic converters, restrictions on movements of heavy vehicles during day time and declaration of low emission zones (Molina and Molina, 2002; CPCB, 2006; Davis, 2008; DEAT, 2008; Molina et al., 2009; CAI-Asia, 2010b; CPCB, 2010b; Jazcilevich et al., 2011; Naiker et al., 2012; Salcedo et al., 2012; Zhong et al., 2013b). In Mexico, implementation of UAQM strategies has helped in reducing the concentration of criteria pollutants i.e., CO, SO₂, NO₂ and Pb, whereas concentrations of tropospheric O₃, PM₁₀ and PM_{2.5} are still at critical levels (Molina et al., 2007; Salcedo et al., 2012). In the recent past, Chavez-Baeza and Sheinbaum-Pardo (2014) have estimated that the implementation of low emitting vehicles and BRT in Mexico City Metropolitan Area can significantly reduce urban air pollution levels. In one of the studies, Stankovic et al. (2012) have evaluated sustainable urban air pollution management approach to tackle traffic related air pollution in the city of Banja Luka, Bosnia and Herzegovina through implementation of automatic traffic control system. Further, Menon-Choudhary and Shukla (2009) have described that such modern technologies, if conjoined with effective greenhouse gas mitigation controls, efficient policy governance with independent environment regulatory authority, can help to build an efficient and effective UAQMP in any developing country including India. Table 5 describes various UAQM strategies undertaken to maintain the acceptable air quality in different countries of the world.

3.8. Public participation

Public participation includes active response from citizens and stakeholders in urban air quality goal setting (Longhurst et al., 1996; DEAT, 2008). Karatzas et al. (2003) have described techniques for effective dissemination of urban air quality information to the public using mobile applications, street panels and mass media. Most of the EU countries are providing air quality information on websites and through publications (DNERI, 2004). In the USA, air quality information is shared with the public via AIRNow (AIRNow, 2011) which provides real time and forecast AQI for 300 cities. In addition to that, the “Window to my Environment” is a web-based tool that provides a wide range of federal, state, and local information about environmental conditions and features of the concerned area within the USA (U.S. EPA, 2011c). In South Africa, the information related to air quality is disseminated through web pages, newsletters and mass media (NILU, 2007). In India and China, the effective use of mass and electronic media, street

panels and web pages has led to sharing the air quality of the urban areas with public (AQCIN, 2014; IITM, 2014).

4. Concluding Remarks

Motorized road transport is the dominant source of urban air pollution in almost all the countries of the world. However, increasing ambient air pollution at urban hot spots is one of the critical problems with frequent violations of NAAQS and/or WHO guidelines for pollutants like PM_{2.5}, PM₁₀ and NO_x. An effective and efficient UAQMP may include all the key components, which may help in sustaining an acceptable ambient air quality. The UAQMPs can be implemented at national, city and/or local levels. In most of the developed countries, the UAQMPs are already being implemented successfully. The UAQMPs like SIP and LAQMP possess efficient communication system between national and local authorities, which ensures its effective implementation and thus maintain the acceptable ambient air quality. These UAQMPs have strict air quality standards/limits for all criteria and hazardous air pollutants; continuous real time air quality monitoring network along with display systems; efficient emission inventory model; air quality modeling and control practices and public participation. In London, congestion and road user charging schemes have been implemented successfully aiming to reduce vehicular pollution in specific defined zones which have significantly reduced CO₂, NO_x and PM₁₀ concentrations by 16.4%, 13.4% and 6.9%, respectively (EEA, 2008). Further, Tonne et al. (2008, 2010) have reported significant reduction in PM and NO_x concentrations after the implementation of “congestion charging” in London which thereafter resulted in an increase in associated health benefits. Hasheminassab et al. (2014) have evaluated the impact of UAQM strategies in the reduction of PM_{2.5} emissions from vehicular source using SA. Results have indicated that PM_{2.5} emissions from the year 2002 to 2012 have been decreased by 24% and 21% in Los Angeles and Rubidoux, respectively. Another successful implementation of UAQMP can be observed at Cardiff and Norwich cities, where significant reductions in NO₂ concentrations have been achieved (Moorcroft and Dore, 2013). In USA, efficient and effective SIP in regions of Connecticut, Georgia, Illinois, Indiana, Kentucky, Maryland, Michigan, Missouri, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Tennessee and West Virginia and the District of Columbia has helped in achieving the goal of bringing down the concentrations of PM_{2.5} within the prescribed standards (Cohan and Chen, 2014). Soret et al. (2011) and Soret et al. (2013) have described how the improvements in urban traffic fleet and vehicle technologies can significantly reduce ambient concentrations of NO₂ and PM₁₀ in the city of Barcelona, Spain. Using WRF-ARW/HERMES/CMAQ modeling system at very high spatial and temporal resolutions, it is observed that NO₂ and PM₁₀ reductions are in the range of 15 µg/m³ and 5 µg/m³, respectively. In addition, Henschel et al. (2012) have reviewed control strategies that have been implemented to reduce air pollution levels and their health impacts. They have reported that interventions have successfully reduced air pollution levels and increased associated health benefits, mainly through reduced cardiovascular and/or respiratory mortality and/or morbidity. Further, Pope et al. (2009) have observed that a reduction in the exposure to ambient fine PM concentrations has resulted in significant and measurable improvements in life expectancy in 211 county units in 51 U.S. metropolitan areas. The decrease of 10 µg/m³ in the concentration of PM_{2.5} has been associated with an increase in mean life expectancy of 0.61 years in urban and densely populated counties. Later, Correia et al. (2013) carried out similar studies in 545 counties in USA and found that reductions in PM_{2.5} concentrations due to implementation of efficient and effective UAQM strategies resulted in an increase in life expectancy. Similarly, in EEA member countries, a decrease in emission levels of NO_x and PM_{2.5} from transport sector from 1990 to 2005 has resulted in a reduction in health impacts in the range of 2.5% in Bulgaria to 25% in Luxembourg and Switzerland (EEA, 2010). In another study, involving 25 cities of 12 European countries, it has been estimated that decrease of long-term exposure to PM_{2.5} to 10 µg/m³ can

prevent more than 19 000 premature deaths, annually (APHEKOM, 2011). In addition, Pererz et al. (2009) have estimated that reducing the mean PM₁₀ exposure by 30 mg/m³ has resulted in 3 500 lesser deaths, 1 800 lesser hospitalizations for cardio-respiratory diseases, 5 100 lesser cases of chronic bronchitis among adults, 31 100 lesser cases of acute bronchitis among children and 54 000 lesser asthma attacks among children and adults. As a consequence, the mean total monetary benefits have been estimated to be 6 400 million Euros per year. Table 6 has described

some examples of success stories of achieving health benefits due to reduction in PM concentrations after implementing UAQM strategies. However, in most of the developing countries, UAQMPs are either in process of development or do not exist at all. Hence, looking at the increasing urbanization globally, the need of the hour is to equip the air quality regulatory authorities with effective and efficient UAQMP, which may help them to maintain the urban quality within the prescribed limits/standards.

Table 5. UAQM strategies of developed and developing countries

UAQM Strategies	US ^a	UK ^b	EU ^c	Australia ^d	British Columbia ^e	Japan ^f	Hong Kong ^g	Singapore ^h	Mexico ⁱ	Thailand ^j	South Africa ^k	China ^l	India ^m
Alternative fuel option in vehicles	x	x	x	x	x	x	x	x	x	x	o	x	x
Improvement in fuel quality	x	x	x	x	x	x	x	x	x	o	o	x	x
Fitting of catalytic converter	x	x	x	x	x	x	x	x	x	o	o	o	o
Inspection and maintenance program	x	x	x	x	x	x	x	x	x	o	o	o	o
Use of low and zero emission vehicles	x	x	x	x	x	x	x	o	x	o	o	o	o
Banned on street parking	x	x	x	x	x	x	x	x	x	o	o	o	o
Support for cycling	x	x	x	x	x	x	x	x	o	x	–	x	n
Congestion charging	x	x	x	x	–	–	–	–	x	n	n	n	n
Specific bus corridors	x	x	x	x	x	x	o	o	x	o	o	o	o
Pedestrians corridors	x	x	x	x	x	x	o	o	x	o	o	o	o
Low emission zone	–	x	x	o	–	–	–	n	o	n	n	n	n
Banned on entry of old vehicles in air quality control regions	x	x	x	x	x	x	x	x	x	o	o	o	o
Banned of heavy good vehicles in air quality control regions	x	x	x	x	x	x	x	x	x	o	o	o	o
Banned on idling of vehicles at traffic signal	x	x	x	o	x	–	o	–	–	–	n	–	n
Split Cycle Offset Optimization Technique (SCOOT) system	o	x	x	–	–	–	n	n	–	n	n	n	n
Intelligent transport system	x	x	x	–	–	–	–	–	–	–	n	n	n
Banned on smoky vehicle	x	x	x	x	x	x	x	x	x	x	o	o	o
On-board diagnostic system in vehicle	x	x	x	x	x	x	x	x	x	x	o	o	o
Subsidy on registration tax on environment-friendly vehicles	x	x	x	–	x	x	–	–	x	n	n	n	n
Mass rapid transport	x	x	x	x	x	x	x	x	x	x	o	x	x
Encouraging carpools	x	x	x	x	x	–	–	x	–	o	n	n	n
Maintenances of road infrastructure	x	x	x	o	x	x	o	x	x	o	o	o	o
Stringent emission norms	x	x	x	x	x	x	x	x	x	o	o	x	x
Reduction in diesel vehicles	x	x	x	x	x	x	x	x	x	o	n	n	n

^a CULS (2014), U.S. EPA (2011a)

^b Corporation of London (2003), Newcastle City Council (2006)

^c EEA (2011b), EEA (2013b), EEA (2013c), EU-Transport and Environment (2014)

^d NSW Government (2009), Commonwealth of Australia (2011), EPA (2007), Todd et al. (2011)

^e B. C. Government (2008)

^f CEC (2005), CEC (2011)

^g Environmental Bureau, Hong Kong (2013), Edesess (2011)

^h ADB (2006a), NEA (2014)

ⁱ Molina and Molina (2002), Davis (2008), Molina et al. (2009), Molina et al. (2010), Jazcilevich et al. (2011), Salcedo et al. (2012), Chavez-Baeza and Sheinbaum-Pardo (2014)

^j KEI (2002); IGES (2007), CAI-Asia (2010b)

^k DEAT (2008), Naiker et al. (2012)

^l Hao et al. (2005), Zhou et al. (2010), Wu et al. (2011), Zhong et al. (2013b)

^m CPCB (2006), CPCB (2010b)

x=properly implemented, o=partially implemented, n=not implemented, – information not available

Table 6. Health benefits of implementation of UAQM strategies

Country	References	Case Study	Main Outcomes
USA	Friedman et al. (2001)	Traffic management during Olympic games period in 1996 at Atlanta	<ul style="list-style-type: none"> Significantly decreased in PM (16.1%) and O₃ (13–30%) concentrations during Olympic period compared to baseline period
	Pope et al. (2009)	Air pollution health impacts result compared with baseline in surrounding years	<ul style="list-style-type: none"> Significant decrease in number of hospital visit by Asthma patients
	Peel et al. (2010)	Short as well as long term management	<ul style="list-style-type: none"> Increase in mean life expectancy of years in 545 county in US due to decrease in PM₁₀ concentrations
	Correia et al. (2013)	Impacts assessment of European Air Emission Policies	<ul style="list-style-type: none"> Air quality and public health improved in all 32 EEA members states
European Environment Agency Members UK	EEA (2010)	Long term traffic management	<ul style="list-style-type: none"> Reduction in PM and NO₂ concentrations
	APHEKOM (2011)	The London congestion charging scheme	<ul style="list-style-type: none"> Decrease in years of life lost in range of 2.5–25% through EEA members
Sweden	Tonne et al. (2008)	Long term traffic management	<ul style="list-style-type: none"> Decrease in NO₂ and PM₁₀ concentration level in designated area
	Tonne et al. (2010)	The Stockholm congestion charging scheme	<ul style="list-style-type: none"> Increase in years life gained of resident live within congestion charging zone
China	Johansson et al. (2009)	Long term traffic management	<ul style="list-style-type: none"> Significant reduction in hospital admission of bronchitis patients
	Huang et al. (2009b)	Traffic management during Olympic games in Beijing in 2008	<ul style="list-style-type: none"> Significant decrease in NO_x (10%) and PM₁₀ (7.6%) concentration within CCZ compared with previous year
	Li et al. (2010)	Short term traffic management	<ul style="list-style-type: none"> 210 year life gained per 100 000 persons from Greater Stockholm over a 10 year period Significant decrease in asthma patients visit to hospitals. Increase in 10 µg/m³ of PM_{2.5} and 10 ppb of O₃ found to be associated with increase in 2%–4.4% in asthma outpatients visits

References

- ADB (Asian Development Bank), 2006a. Country Synthesis Report on Urban Air Quality Management for Singapore, Asian Development Bank and the Clean Air Initiative for Asian Cities (CAI-Asia) Centre, Philippines, 22 pages.
- ADB (Asian Development Bank), 2006b. Country Synthesis Report on Urban Air Quality Management for India, Asian Development Bank and the Clean Air Initiative for Asian Cities (CAI-Asia) Center, 34 pages.
- Agarwal, R., Jayaraman, G., Anand, S., Marimuthu, P., 2006. Assessing respiratory morbidity through pollution status and meteorological conditions for Delhi. *Environmental Monitoring and Assessment* 114, 489–504.
- Air quality-BC, 2014. www.envistaweb.env.gov.bc.ca, accessed in May 2014.
- Air quality-China, 2014. www.aqicn.org, accessed in May 2014.
- Air quality-EU, 2014. www.eea.europa.eu/data-and-maps/data/airbase, accessed in May 2014.
- Air quality-India, 2014. <http://164.100.43.188/cpcbnew>, accessed in January 2014.
- Air quality-NSW, 2014. www.environment.nsw.gov.au/aqms, accessed in April, 2014.
- Air quality-Mexico, 2014. http://sinaica2.inecc.gob.mx/magic/rmart/null/_framework, accessed in May 2014.
- Air quality-South Africa, 2014. www.saaqis.org.za, accessed in April 2014.
- Air quality-USA, 2014. www.airnow.gov, accessed in June, 2014.
- Air quality-UK, 2014. www.uk-air.defra.gov.uk, accessed in May, 2014.
- AIRNow, 2011. http://www.airnow.gov/index.cfm?action=topics.about_airnow, accessed in November 2011.
- Almeida, S.M., Pio, C.A., Freitas, M.C., Reis, M.A., Trancoso, M.A., 2006. Source apportionment of atmospheric urban aerosol based on weekdays/weekend variability: Evaluation of road re-suspended dust contribution. *Atmospheric Environment* 40, 2058–2067.
- Amato, F., Cassee, F.R., van der Gon, H.A.C.D., Gehrig, R., Gustafsson, M., Hafner, W., Harrison, R.M., Jozwicka, M., Kelly, F.J., Moreno, T., Prevot, A.S.H., Schaap, M., Sunyer, J., Querol, X., 2014. Urban air quality: The challenge of traffic non-exhaust emissions. *Journal of Hazardous Materials* 275, 31–36.
- Anjaneyulu, M.V.L.R., Harikrishna, M., Chenchubulu, S., 2006. Modeling ambient carbon monoxide pollutant due to road traffic. *Proceedings of World Academy of Science, Engineering and Technology* 17, 103–106.
- APHEKOM, 2011. Improving Knowledge and Communication for Decision Making on Air Pollution and Health in Europe, Summary Report of the APHEKOM Project 2008–2011, 12 pages.
- AQCC (Air Quality Coordinating Committee), 2012. Perth Air Quality Management Plan Report Card 2010/11, Department of Environment and Conservation, Government of Western Australia, 28 pages.
- AQCIN, 2014. <http://aqicn.org/city>, accessed in January, 2014.
- AQS-Australia (Air quality standards), 2014. <http://www.environment.gov.au/topics/environment-protection/air-quality/air-quality-standards#air>, accessed in May 2014.
- AQS-BC (Air quality standards for British Columbia), 2014. <http://www.bcairquality.ca/reports/pdfs/aqotable.pdf>, accessed in April 2014.
- AQS-China (Air quality standards for China), 2014. http://www.vecc-sepa.org.cn/eng/news/news_detail.jsp?newsid=e00397, accessed in March 2014.
- AQS-Mexico (Air quality standards for Mexico), 2014. http://transportpolicy.net/index.php?title=Mexico:_Air_Quality_Standards, accessed in April 2014.
- ARAI (Automotive Research Association of India), 2007. Emission Factor Development for Indian Vehicles, Project Report No. AEF/2006–07/IOCL/Emission Factor Project, Automotive Research Association of India, Pune, India, 94 pages.
- Atash, F., 2007. The deterioration of urban environments in developing countries: Mitigating the air pollution crisis in Tehran, Iran. *Cities* 24, 399–409.
- Auto Fuel Policy, 2002. Report of the Expert Committee on Auto Fuel Policy – Executive Summary, Ministry of Petroleum and Natural Gases, Government of India, 48 pages.
- Badami, M.G., 2005. Transport and urban air pollution in India. *Environmental Management* 36, 195–204.
- B.C. Government, 2008. A Report on B.C. Air Action Plan, Government of British Columbia, 28 pages.
- Balakrishnan, K., Ganguli, B., Ghosh, S., Sankar, S., Thanasekaran, V., Rayudu, V.N., Caussy, H., 2011. Part 1. Short-term effects of air pollution on mortality: Results from a time-series analysis in Chennai, India. *Research Report – Health Effects Institute* 157, 7–44.
- Baldasano, J.M., Valera, E., Jimenez, P., 2003. Air quality data from large cities. *Science of the Total Environment* 307, 141–165.
- Beattie, C.I., Longhurst, J.W.S., Woodfield, N.K., 2001. Air quality management: Evolution of policy and practice in the UK as exemplified by the experience of English local government. *Atmospheric Environment* 35, 1479–1490.
- Begum, B.A., Biswas, S.K., Hopke, P.K., 2011. Key issues in controlling air pollutants in Dhaka, Bangladesh. *Atmospheric Environment* 45, 7705–7713.
- Begum, B.A., Biswas, S.K., Markwitz, A., Hopke, P.K., 2010. Identification of sources of fine and coarse particulate matter in Dhaka, Bangladesh. *Aerosol and Air Quality Research* 10, 345–353.
- Belis, C.A., Karagulian, F., Larsen, B.R., Hopke, P.K., 2013. Critical review and meta-analysis of ambient particulate matter source apportionment using receptor models in Europe. *Atmospheric Environment* 69, 94–108.
- Benson, P.E., 1984. CALINE4–A Dispersion Model for Predicting Air Pollutant Concentrations Near Roadways, FHWA–CA–TL–84–15, California Department of Transportation, Sacramento, CA, 247 pages.
- Blifford Jr, I.H., Meeker, G.O., 1967. A factor analysis model of large scale pollution. *Atmospheric Environment (1967)* 1, 147–157.
- Boulter, P., McCrae, I., Barlow, T., 2007. A Review of Instantaneous Emissions Models for Road Vehicles, Transport Research Laboratory, Research Report 267, Transport Research Laboratory, Wokingham, UK, 55 pages.
- Bove, M.C., Brotto, P., Cassola, F., Cuccia, E., Massabo, D., Mazzino, A., Piazzalunga, A., Prati, P., 2014. An integrated PM_{2.5} source apportionment study: Positive Matrix Factorisation vs. The chemical transport model CAMx. *Atmospheric Environment* 94, 274–286.
- Breitner, S., Liu, L.Q., Cyrus, J., Bruske, I., Franck, U., Schlink, U., Leitte, A.M., Herbarth, O., Wiedensohler, A., Wehner, B., Hu, M., Pan, X.C., Wichmann, H.E., Peters, A., 2011. Sub-micrometer particulate air pollution and cardiovascular mortality in Beijing, China. *Science of the Total Environment* 409, 5196–5204.
- BTRE (Bureau of Transport and Regional Economics), 2005. Health Impacts of Transport Emissions in Australia: Economic Costs, Working Paper 63, Commonwealth of Australia, 169 pages.
- Byun, D., Schere, K.L., 2006. Review of the governing equations, computational algorithms, and other components of the models–3 Community Multiscale Air Quality (CMAQ) modeling system. *Applied Mechanics Reviews* 59, 51–77.
- CAAC (Clean Air Alliance of China), 2013. “Twelfth Five-Year Plan” on Air Pollution Prevention and Control in Key Regions, Clean Air Alliance of China, Ministry of Environmental Protection, China, 76 pages.
- CAI-Asia (Clean Air Initiative for Asian Cities), 2010a. Air Quality in Asia: Status and Trends, 2010 Edition, CAI-Asia Center, Philippines, 25 pages.
- CAI-Asia (Clean Air Initiative for Asian Cities), 2010b. Thailand Air Quality Profile: 2010 Edition, Draft Report, CAI-Asia Center, Philippines, 39 pages.

- Carruthers, D.J., Edmunds, H.A., Lester, A.E., McHugh, C.A., Singles, R.J., 2000. Use and validation of ADMS–Urban in contrasting urban and industrial locations. *International Journal of Environment and Pollution* 14, 364–374.
- CEC (Central Environmental Council), 2011. Future Policy for Motor Vehicle Emission Reduction (10th Report), Central Environmental Council, Government of Japan, 2 pages.
- CEC (Central Environmental Council), 2005. Future Policy for Motor Vehicle Emission Reduction (8th Report), Central Environmental Council, Government of Japan, 66 pages.
- Chakrobarty, A., Gupta, T., 2010. Chemical characterization of submicron aerosol in Kanpur region: a source apportionment study. *Aerosol and Air Quality Research* 10, 433–445.
- Chan, C.K., Yao, X., 2008. Air pollution in mega cities in China. *Atmospheric Environment* 42, 1–42.
- Chavez–Baeza, C., Sheinbaum–Pardo, C., 2014. Sustainable passenger road transport scenarios to reduce fuel consumption, air pollutants and GHG (greenhouse gas) emissions in the Mexico City Metropolitan Area. *Energy* 66, 624–634.
- Chelani, A.B., Gajghate, D.G., Devotta, S., 2008. Source apportionment of PM₁₀ in Mumbai, India using CMB model. *Bulletin of Environmental Contamination and Toxicology* 81, 190–195.
- Cheung, K., Daher, N., Kam, W., Shafer, M.M., Ning, Z., Schauer, J.J., Sioutas, C., 2011. Spatial and temporal variation of chemical composition and mass closure of ambient coarse particulate matter (PM_{10–2.5}) in the Los Angeles area. *Atmospheric Environment* 45, 2651–2662.
- Chhabra, S.K., Chhabra, P., Rajpal, S., Gupta, R.K., 2001. Ambient air pollution and chronic respiratory morbidity in Delhi. *Archives of Environmental Health* 56, 58–64.
- Chow, J.C., Watson, J.G., Lowenthal, D.H., Chen, L.W.A., Motallebi, N., 2011. PM_{2.5} source profiles for black and organic carbon emission inventories. *Atmospheric Environment* 45, 5407–5414.
- Chowdhury, Z., Zheng, M., Schauer, J.J., Sheesley, R.J., Salmon, L.G., Cass, G.R., Russell, A.G., 2007. Speciation of ambient fine organic carbon particles and source apportionment of PM_{2.5} in Indian cities. *Journal of Geophysical Research–Atmospheres* 112, art. no. D15303.
- Cimorelli, A.J., Perry, S.G., Venkatram, A., Weil, J.C., Paine, R.J., Wilson, R.B., Lee, R.F., Peters, W.D., Brode, R.W., 2005. AERMOD: A dispersion model for industrial source applications. Part I: General model formulation and boundary layer characterization. *Journal of Applied Meteorology* 44, 682–693.
- Cities Alliance, 2007. Liveable Cities, The Benefits of Urban Environmental Planning, A Cities Alliance Study on Good Practices and Useful Tools, Washington, D.C., USA, 162 pages.
- Cohan, D.S., Chen, R., 2014. Modeled and observed fine particulate matter reductions from state attainment demonstrations. *Journal of the Air & Waste Management Association* 64, 995–1002.
- Colais, P., Faustini, A., Stafoggia, M., Berti, G., Bisanti, L., Cadum, E., Cernigliaro, A., Mallone, S., Pacelli, B., Serinelli, M., Simonato, L., Vigotti, M.A., Forastiere, F., Grp, E.C., 2012. Particulate air pollution and hospital admissions for cardiac diseases in potentially sensitive subgroups. *Epidemiology* 23, 473–481.
- Colville, R.N., Gomez–Perales, J.E., Nieuwenhuijsen, M.J., 2003. Use of dispersion modelling to assess road–user exposure to PM_{2.5} and its source apportionment. *Atmospheric Environment* 37, 2773–2782.
- Commonwealth of Australia, 2011. State of the Environment, Australia, Available at national library of Australia, ISBN 978–1–921733–35–2, 940 pages.
- Cooper, J.A., Watson, J.G., 1980. Receptor oriented methods of air particulate source apportionment. *Journal of the Air Pollution Control Association* 30, 1116–1125.
- Corporation of London, 2003. Air Quality Action Plan, The Corporation of London, UK., 58 pages.
- Correia, A.W., Pope, C.A., Dockery, D.W., Wang, Y., Ezzati, M., Dominici, F., 2013. Effect of air pollution control on life expectancy in the United States an analysis of 545 US counties for the period from 2000 to 2007. *Epidemiology* 24, 23–31.
- Coutant, B.W., Engel–Cox, J., Swinton, K.E., 2003. Compilation of Existing Studies on Source Apportionment for PM_{2.5}, Technical Report, Columbus, Ohio, 88 pages.
- CPCB (Central Pollution Control Board), 2014. <http://cpcb.nic.in/air.php>, accessed in January 2014.
- CPCB (Central Pollution Control Board), 2010a. Air Quality Monitoring, Emission Inventory and Source Apportionment Study for Indian Cities, National Summary Report. The Central Pollution Control Board, New Delhi, India, 290 pages.
- CPCB (Central Pollution Control Board), 2010b. Status of the Vehicular Pollution Control Programme in India, The Central Pollution Control Board, New Delhi, India, Probes/136, 114 pages.
- CPCB (Central Pollution Control Board), 2006. Air Quality Trends and Action Plan for Control of Air Pollution from Seventeen Cities, Ministry of Environment and Forest, NAAQMS/29/2006–07, New Delhi, India, 218 pages.
- CPCB (Central Pollution Control Board), 1998. Assessment of Impact to Air Environment: Guidelines for Conducting Air Quality Modelling, Privesh Newsletter, The Central Pollution Control Board, New Delhi.
- CULS (Cornell University Law School), 2014. <http://www.law.cornell.edu/uscode/text/42/7408>, accessed in June 2014.
- Daher, N., Hasheminassab, S., Shafer, M.M., Schauer, J.J., Sioutas, C., 2013. Seasonal and spatial variability in chemical composition and mass closure of ambient ultrafine particles in the megacity of Los Angeles. *Environmental Science–Processes & Impacts* 15, 283–295.
- Davis, L.W., 2008. The effect of driving restrictions on air quality in Mexico City. *Journal of Political Economy* 116, 38–81.
- DEA (Department of Environmental Affairs), 2011. Baseline Survey on Government owned Air Quality Monitoring Networks Report, Prepared by Atmospheric Quality Information Directorate, Department of Environmental Affairs, Republic of South Africa, 31 pages.
- DEA (Department of Environmental Affairs), 2009. National Ambient Air Quality Standards, National Environmental Management: Air Quality Act, 2004, Government Gazette, Vol. 534, No. 32816, Republic of South Africa.
- DEAT (Department of Environmental Affairs and Tourism), 2008. Manual for Air Quality Management Planning, Department of Environmental Affairs and Tourism, Republic of South Africa, 95 pages.
- DEFRA (Department of Environment Federation and Rural Affairs), 2013. <http://uk-air.defra.gov.uk/networks>, accessed in December, 2013.
- DEFRA (Department of Environment Federation and Rural Affairs), 2011a. <http://uk-air.defra.gov.uk/air-pollution/uk-eu-limits>, accessed in December 2011.
- DEFRA (Department of Environment Federation and Rural Affairs), 2011b. <http://uk-air.defra.gov.uk/networks/network-info?view=aurng>, accessed in December 2011.
- DEFRA (Department of Environment Federation and Rural Affairs), 2011c. <http://naei.defra.gov.uk>, accessed in December, 2011.
- DEFRA (Department of Environment Federation and Rural Affairs), 2009a. Local Air Quality Management Policy Guidance (PG09), Department of Environment Federation and Rural Affairs, Government of United Kingdom, 47 pages.
- DEFRA (Department of Environment Federation and Rural Affairs), 2009b. Local Air Quality Management, Technical Guidance (LAQM–TG09) under Part IV of the Environment Act 1995, Environment (Northern Ireland) Order 2002 Part III, 318 pages.
- DEFRA (Department of Environment Federation and Rural Affairs), 1998. Local Air Quality Management, Technical Guidance (LAQM–TG) under Part IV of the Environment Act 1995, Environment (Northern Ireland) Order 2002 Part III.

- DNERI (The Danish Environment Research Institute), 2004. Comparison of the EU and US Air Quality Standards and Planning Requirements, A project of DG environment Carried out by Milieu Ltd., The Danish Environment Research Institute and The Center for Clean Air Policy, 58 pages.
- Dockery, D.W., 2009. Health effects of particulate air pollution. *Annals of Epidemiology* 19, 257–263.
- Dockery, D.W., Pope, C.A., 3rd, Xu, X., Spengler, J.D., Ware, J.H., Fay, M.E., Ferris, B.G., Jr., Speizer, F.E., 1993. An association between air pollution and mortality in six U.S. cities. *New England Journal of Medicine* 329, 1753–1759.
- Duan, F.K., He, K.B., Ma, Y.L., Yang, F.M., Yu, X.C., Cadle, S.H., Chan, T., Mulawa, P.A., 2006. Concentration and chemical characteristics of PM_{2.5} in Beijing, China: 2001–2002. *Science of the Total Environment* 355, 264–275.
- Duffield, R., 2011. City of London Air Quality Strategy 2011–2015, Port Health and Environmental Services Committee, Corporation of London, London, UK, 75 pages.
- Edesess, M., 2011. Roadside Air Pollution in Hong Kong: Why is It still so Bad?, School of Energy and Environment, City University of Hong Kong, 19 pages.
- EEA (European Environment Agency), 2014. <http://acm.eionet.europa.eu>, accessed in January 2014.
- EEA (European Environment Agency), 2013a. Air Quality in Europe — 2013, EEA Technical Reports 9/2013, Copenhagen, 112 pages.
- EEA (European Environment Agency), 2013b. Road User Charges for Heavy Goods Vehicles (HGV), EEA Technical Report No.1/2013, Copenhagen, 88 pages.
- EEA (European Environment Agency), 2013c. A Closure Look at Urban Transport, Transport Indicators Tracking Progress towards Environmental Targets in Europe, EEA Technical Report No.11/ 2013, Copenhagen, 112 pages.
- EEA (European Environment Agency), 2011a. Air Quality in Europe — 2011 Report, EEA Technical Report No.12/2011, Copenhagen, 88 pages.
- EEA (European Environment Agency), 2011b. Laying the Foundations for Greener Transport: Transport Indicators Tracking Progress towards Environmental Targets in Europe, Copenhagen, EEA Technical Report No.7/2011, 92 pages.
- EEA (European Environment Agency), 2010. Impact of Selected Policy Measures on Europe's Air Quality, Report No. 8, European Environment Agency, Copenhagen, 70 pages.
- EEA (European Environmental Agency), 2008. Success Stories within the Road Transport Sector on Reducing Greenhouse Gas Emission and Producing Ancillary Benefits, EEA Technical Report No. 2/2008, Copenhagen, 70 pages.
- Ekstrom, M., Sjodin, A., Andreasson, K., 2004. Evaluation of the COPERT III emission model with on-road optical remote sensing measurements. *Atmospheric Environment* 38, 6631–6641.
- Elbir, T., 2004. A GIS based decision support system for estimation, visualization and analysis of air pollution for large Turkish cities. *Atmospheric Environment* 38, 4509–4517.
- Elbir, T., Mangir, N., Kara, M., Simsir, S., Eren, T., Ozdemir, S., 2010. Development of a GIS-based decision support system for urban air quality management in the city of Istanbul. *Atmospheric Environment* 44, 441–454.
- Elsom, D., 1996. Smog Alert—Managing Urban Air Quality, Routledge, 192 pages.
- Environmental Bureau, Hong Kong, 2013. A Clean Air Plan for Hong Kong, A report Prepared in Collaboration with Transport and Housing Bureau, Food and Health Bureau and Development Bureau, Hong Kong, 43 pages.
- EPA (Environmental Protection Authority), 2007. Perth Air Quality Management Plan: Five-Year Review. Review Steering Committee report, Western Australia, 52 pages.
- EPD (Environment Protection Department), 2011. Air Quality in Hong Kong, The Government of the Hong Kong, Special Administrative Region, Report No. EPD/TR2/12, 72 pages.
- Esteves-Booth, A., Muneer, T., Kubie, J., Kirby, H., 2002. A review of vehicular emission models and driving cycles. *Proceedings of the Institution of Mechanical Engineers Part C—Journal of Mechanical Engineering Science* 216, 777–797.
- EU-Transport and Environment, 2014. <http://ec.europa.eu/environment/air/transport/road.htm>, accessed in June 2014.
- Faulkner, M., Russell, P., 2010. Review of Local Air Quality Management, A report to DEFRA and The Devolved Administrations, 98 pages.
- Fedra, K., Haurie, A., 1999. A decision support system for air quality management combining GIS and optimization techniques. *International Journal of Environment and Pollution* 12, 125–146.
- Figueiredo, M.L., Monteiro, A., Lopes, M., Ferreira, J., Borrego, C., 2013. Air quality assessment of Estarreja, an urban industrialized area, in a coastal region of Portugal. *Environmental Monitoring and Assessment* 185, 5847–5860.
- Friedman, M.S., Powell, K.E., Hutwagner, L., Graham, L.M., Teague, W.G., 2001. Impact of changes in transportation and commuting behaviors during the 1996 Summer Olympic Games in Atlanta on air quality and childhood asthma. *JAMA—Journal of the American Medical Association* 285, 897–905.
- Gehring, U., Casas, M., Brunekreef, B., Bergstrom, A., Bonde, J.P., Botton, J., Chevrier, C., Cordier, S., Heinrich, J., Hohmann, C., Keil, T., Sunyer, J., Tischer, C.G., Toft, G., Wickman, M., Vrijheid, M., Nieuwenhuijsen, M., 2013. Environmental exposure assessment in European birth cohorts: Results from the ENRIECO project. *Environmental Health* 12, art. no. 8.
- Gietl, J.K., Klemm, O., 2009. Source identification of size-segregated aerosol in Munster, Germany, by factor analysis. *Aerosol Science and Technology* 43, 828–837.
- Gokhale, S., Khare, M., 2007. A theoretical framework for the episodic–urban air quality management plan (e-UAQMP). *Atmospheric Environment* 41, 7887–7894.
- Gokhale, S., Khare, M., 2005. A hybrid model for predicting carbon monoxide from vehicular exhausts in urban environments. *Atmospheric Environment* 39, 4025–4040.
- Graves, R.J., Lee, T.D., McGinnis, L.F., Jr., 1981. Air monitoring network design: A case study. *Journal of Environmental Engineering Division* 107, 941–955.
- Green, G.A., Cummins, P., 2007. Recommendations to the Clean Air Act Advisory Committee, Air Quality Management Subcommittee Phase-II recommendations, United States of America, 99 pages.
- Green, M.C., Chow, J.C., Chang, M.C.O., Chen, L.W.A., Kuhns, H.D., Etyemezian, V.R., Watson, J.G., 2013. Source apportionment of atmospheric particulate carbon in Las Vegas, Nevada, USA. *Particology* 11, 110–118.
- Gu, J.W., Pitz, M., Schnelle-Kreis, J., Diemer, J., Reller, A., Zimmermann, R., Soentgen, J., Stoelzel, M., Wichmann, H.E., Peters, A., Cyrys, J., 2011. Source apportionment of ambient particles: Comparison of positive matrix factorization analysis applied to particle size distribution and chemical composition data. *Atmospheric Environment* 45, 1849–1857.
- Gu, J.X., Bai, Z.P., Liu, A.X., Wu, L.P., Xie, Y.Y., Li, W.F., Dong, H.Y., Zhang, X., 2010. Characterization of atmospheric organic carbon and element carbon of PM_{2.5} and PM₁₀ at Tianjin, China. *Aerosol and Air Quality Research* 10, 167–176.
- Gulliver, J., Briggs, D., 2011. STEMS–Air: A simple GIS-based air pollution dispersion model for city-wide exposure assessment. *Science of the Total Environment* 409, 2419–2429.
- Gummeneni, S., Bin Yusup, Y., Chavali, M., Samadi, S.Z., 2011. Source apportionment of particulate matter in the ambient air of Hyderabad city, India. *Atmospheric Research* 101, 752–764.
- Gupta, I., Kumar, R., 2006. Trends of particulate matter in four cities in India. *Atmospheric Environment* 40, 2552–2566.

- Gupta, I., Salunkhe, A., Kumar, R., 2010. Modelling 10-year trends of PM₁₀ and related toxic heavy metal concentrations in four cities in India. *Journal of Hazardous Materials* 179, 1084–1095.
- Gupta, A.K., Karar, K., Srivastava, A., 2007. Chemical mass balance source apportionment of PM₁₀ and TSP in residential and industrial sites of an urban region of Kolkata, India. *Journal of Hazardous Materials* 142, 279–287.
- Gurjar, B.R., Jain, A., Sharma, A., Agarwal, A., Gupta, P., Nagpure, A.S., Lelieveld, J., 2010. Human health risks in megacities due to air pollution. *Atmospheric Environment* 44, 4606–4613.
- Gurjar, B.R., Butler, T.M., Lawrence, M.G., Lelieveld, J., 2008. Evaluation of emissions and air quality in megacities. *Atmospheric Environment* 42, 1593–1606.
- Gurjar, B.R., van Aardenne, J.A., Lelieveld, J., Mohan, M., 2004. Emission estimates and trends (1990–2000) for megacity Delhi and implications. *Atmospheric Environment* 38, 5663–5681.
- Guttikunda, S.K., Calori, G., 2013. A GIS based emissions inventory at 1 km x 1 km spatial resolution for air pollution analysis in Delhi, India. *Atmospheric Environment* 67, 101–111.
- Han, X.L., Naeher, L.P., 2006. A review of traffic-related air pollution exposure assessment studies in the developing world. *Environment International* 32, 106–120.
- Handscombe, C.M., Elsom, D.M., 1982. Rationalisation of the national survey of air pollution monitoring network of the United Kingdom using spatial correlation analysis: A case-study of the Greater London area. *Atmospheric Environment (1967)* 16, 1061–1070.
- Hao, J.M., Wang, L.T., 2005. Improving urban air quality in China: Beijing case study. *Journal of the Air & Waste Management Association* 55, 1298–1305.
- Hao, J.M., He, D.Q., Wu, Y., Fu, L., He, K.B., 2000. A study of the emission and concentration distribution of vehicular pollutants in the urban area of Beijing. *Atmospheric Environment* 34, 453–465.
- Hasheminassab, S., Daher, N., Ostro, B.D., Sioutas, C., 2014. Long-term source apportionment of ambient fine particulate matter (PM_{2.5}) in the Los Angeles Basin: A focus on emissions reduction from vehicular sources. *Environmental Pollution* 193, 54–64.
- Hasheminassab, S., Daher, N., Schauer, J.J., Sioutas, C., 2013. Source apportionment and organic compound characterization of ambient ultrafine particulate matter (PM) in the Los Angeles Basin. *Atmospheric Environment* 79, 529–539.
- HEI (Health Effects Institute), 2010. Public Health and Air Pollution in Asia (PAPA): Coordinated Studies of Short-Term Exposure to Air Pollution and Daily Mortality in Four Cities, Research Report 154, Health Effects Institute, Boston, MA.
- HEI (Health Effects Institute), 2004. Health Effects of Outdoor Air Pollution in Developing Countries of Asia: A Literature Review, Special Report 15, Health Effects Institute, Boston, MA.
- Henry, R.C., Lewis, C.W., Hopke, P.K., Williamson, H.J., 1984. Review of receptor model fundamentals. *Atmospheric Environment* 18, 1507–1515.
- Henschel, S., Atkinson, R., Zeka, A., Le Tertre, A., Analitis, A., Katsouyanni, K., Chanel, O., Pascal, M., Forsberg, B., Medina, S., Goodman, P.G., 2012. Air pollution interventions and their impact on public health. *International Journal of Public Health* 57, 757–768.
- Herlekar, M., Joseph, A.E., Kumar, R., Gupta, I., 2012. Chemical speciation and source assignment of particulate (PM₁₀) phase molecular markers in Mumbai. *Aerosol and Air Quality Research* 12, 1247–1260.
- Hopke, P.K., 1991. *Receptor Modeling for Air Quality Management*, Elsevier, Amsterdam, the Netherlands, pp. 1–8.
- Huang, W., Tan, J.G., Kan, H.D., Zhao, N., Song, W.M., Song, G.X., Chen, G.H., Jiang, L.L., Jiang, C., Chen, R.J., Chen, B.H., 2009. Visibility, air quality and daily mortality in Shanghai, China. *Science of the Total Environment* 407, 3295–3300.
- Huo, H., Zhang, Q., He, K.B., Wang, Q.D., Yao, Z.L., Streets, D.G., 2009. High-resolution vehicular emission inventory using a link-based method: A case study of light-duty vehicles in Beijing. *Environmental Science & Technology* 43, 2394–2399.
- Hwang, I., Hopke, P.K., Pinto, J.P., 2008. Source apportionment and spatial distributions of coarse particles during the Regional Air Pollution Study. *Environmental Science & Technology* 42, 3524–3530.
- IGES (Institute of Global Environmental Strategies), 2007. Air Pollution Control in the Transport Sector: Third Phase Research Report of the Urban Environmental Project, Japan, 392 pages.
- IITM (Indian Institute of Tropical Meteorology), 2014. <http://safar.tropmet.res.in>, accessed in January 2014.
- Ito, K., Xue, N., Thurston, G., 2004. Spatial variation of PM_{2.5} chemical species and source-apportioned mass concentrations in New York City. *Atmospheric Environment* 38, 5269–5282.
- Janssen, N.A.H., Schwartz, J., Zanobetti, A., Suh, H.H., 2002. Air conditioning and source-specific particles as modifiers of the effect of PM₁₀ on hospital admissions for heart and lung disease. *Environmental Health Perspectives* 110, 43–49.
- Jazcilevich, A.D., Reynoso, A.G., Grutter, M., Delgado, J., Ayala, U.D., Lastra, M.S., Zuk, M., Oropeza, R.G., Lents, J., Davis, N., 2011. An evaluation of the hybrid car technology for the Mexico Mega City. *Journal of Power Sources* 196, 5704–5718.
- Johansson, C., Burman, L., Forsberg, B., 2009. The effects of congestions tax on air quality and health. *Atmospheric Environment* 43, 4843–4854.
- Kandlikar, M., 2007. Air pollution at a hotspot location in Delhi: Detecting trends, seasonal cycles and oscillations. *Atmospheric Environment* 41, 5934–5947.
- Kar, S., Maity, J.P., Samal, A.C., Santra, S.C., 2010. Metallic components of traffic-induced urban aerosol, their spatial variation, and source apportionment. *Environmental Monitoring and Assessment* 168, 561–574.
- Karatzas, K.D., 2002. Theoretical investigation of urban air quality management systems performance towards simplified strategic environmental planning. *Water, Air and Soil Pollution: Focus* 2, 669–676.
- Karatzas, K., Dioudi, E., Moussiopoulos, N., 2003. Identification of major components for integrated urban air quality management and information systems via user requirements prioritisation. *Environmental Modelling & Software* 18, 173–178.
- KEI (Korea Environment Institute), 2002. Benchmarking Urban air Quality Management and Practice in Mega and Major Cities in Asia, Air Pollution in the Mega Cities of Asia (APMA) Project, 129 pages.
- Keller, M., Zbinden, R., 2004. Pollutant Emissions from Road Transport 1980–2030, Swiss Agency for the Environment, Forests and Landscape (SAEFL), Environmental Series No. 355, Berne, Switzerland, 102 pages.
- Kertesz, Z., Szoboszlai, Z., Angyal, A., Dobos, E., Borbely-Kiss, I., 2010. Identification and characterization of fine and coarse particulate matter sources in a middle-European urban environment. *Nuclear Instruments & Methods in Physics Research Section B-Beam Interactions with Materials and Atoms* 268, 1924–1928.
- Kesarkar, A.P., Dalvi, M., Kaginalkar, A., Ojha, A., 2007. Coupling of the Weather Research and Forecasting Model with AERMOD for pollutant dispersion modeling. A case study for PM₁₀ dispersion over Pune, India. *Atmospheric Environment* 41, 1976–1988.
- Khare, P., Baruah, B.P., 2010. Elemental characterization and source identification of PM_{2.5} using multivariate analysis at the suburban site of North-East India. *Atmospheric Research* 98, 148–162.
- Khare, M., Sharma, P., 1999. Performance evaluation of general finite line source model for Delhi traffic conditions. *Transportation Research Part D-Transport and Environment* 4, 65–70.
- Khare, M., Nagendra, S., Gulia, S., 2012. Performance evaluation of air quality dispersion models at urban intersection of an Indian city: a case study of Delhi city. *Proceeding of 20th WIT International Conference on Air Pollution*, May 16–18, 2012, Coruna, Spain, pp. 249–259.

- Khillare, P.S., Balachandran, S., Meena, B., 2004. Spatial and temporal variation of heavy metals in atmospheric aerosol of Delhi. *Environmental Monitoring and Assessment* 90, 1–21.
- Kong, S.F., Han, B., Bai, Z.P., Chen, L., Shi, J.W., Xu, Z., 2010. Receptor modeling of PM_{2.5}, PM₁₀ and TSP in different seasons and long-range transport analysis at a coastal site of Tianjin, China. *Science of the Total Environment* 408, 4681–4694.
- Kota, S.H., Zhang, H.L., Chen, G., Schade, G.W., Ying, Q., 2014. Evaluation of on-road vehicle CO and NO_x National Emission Inventories using an urban-scale source-oriented air quality model. *Atmospheric Environment* 85, 99–108.
- Kothai, P., Saradhi, I.V., Prathibha, P., Hopke, P.K., Pandit, G.G., Puranik, V.D., 2008. Source apportionment of coarse and fine particulate matter at Navi Mumbai, India. *Aerosol and Air Quality Research* 8, 423–436.
- Kuo, C.P., Liao, H.T., Chou, C.C.K., Wu, C.F., 2014. Source apportionment of particulate matter and selected volatile organic compounds with multiple time resolution data. *Science of the Total Environment* 472, 880–887.
- Kura, B., Verma, S., Ajdari, E., Iyer, A., 2013. Growing public health concerns from poor urban air quality: Strategies for sustainable urban living. *Computational Water, Energy, and Environmental Engineering* 2, 1–9.
- Laden, F., Neas, L.M., Dockery, D.W., Schwartz, J., 2000. Association of fine particulate matter from different sources with daily mortality in six US cities. *Environmental Health Perspectives* 108, 941–947.
- Laupsa, H., Denby, B., Larssen, S., Schaugh, J., 2009. Source apportionment of particulate matter (PM_{2.5}) in an urban area using dispersion, receptor and inverse modelling. *Atmospheric Environment* 43, 4733–4744.
- Laxen D., 1993. An introduction to local air quality management. A supplement to *Clean Air* 23, 12.
- Leiman, A., Standish, B., Boting, A., van Zyl, H., 2007. Reducing the healthcare costs of urban air pollution: The South African experience. *Journal of Environmental Management* 84, 27–37.
- Leitte, A.M., Schlink, U., Herbarth, O., Wiedensohler, A., Pan, X.C., Hu, M., Richter, M., Wehner, B., Tuch, T., Wu, Z.J., Yang, M.J., Liu, L.Q., Breitner, S., Cyrys, J., Peters, A., Wichmann, H.E., Franck, U., 2011. Size-segregated particle number concentrations and respiratory emergency room visits in Beijing, China. *Environmental Health Perspectives* 119, 508–513.
- Li, J.J., Wang, G.H., Aggarwal, S.G., Huang, Y., Ren, Y.Q., Zhou, B.H., Singh, K., Gupta, P.K., Cao, J.J., Zhang, R., 2014. Comparison of abundances, compositions and sources of elements, inorganic ions and organic compounds in atmospheric aerosols from Xi'an and New Delhi, two megacities in China and India. *Science of the Total Environment* 476, 485–495.
- Li, G., Zavala, M., Lei, W., Tsimpidi, A.P., Karydis, V.A., Pandis, S.N., Canagaratna, M.R., Molina, L.T., 2011. Simulations of organic aerosol concentrations in Mexico City using the WRF-CHEM model during the MCMA-2006/MILAGRO campaign. *Atmospheric Chemistry and Physics* 11, 3789–3809.
- Li, Y., Wang, W., Kan, H.D., Xu, X.H., Chen, B.H., 2010. Air quality and outpatient visits for asthma in adults during the 2008 Summer Olympic Games in Beijing. *Science of the Total Environment* 408, 1226–1227.
- Lin, L., Lee, M.L., Eatough, D.J., 2010. Review of recent advances in detection of organic markers in fine particulate matter and their use for source apportionment. *Journal of the Air & Waste Management Association* 60, 3–25.
- Lipfert, F.W., Wyzga, R.E., Baty, J.D., Miller, J.P., 2006. Traffic density as a surrogate measure of environmental exposures in studies of air pollution health effects: Long-term mortality in a cohort of US veterans. *Atmospheric Environment* 40, 154–169.
- Liu, L.Q., Breitner, S., Schneider, A., Cyrys, J., Bruske, I., Franck, U., Schlink, U., Leitte, A.M., Herbarth, O., Wiedensohler, A., Wehner, B., Pan, X.C., Wichmann, H.E., Peters, A., 2013. Size-fractionated particulate air pollution and cardiovascular emergency room visits in Beijing, China. *Environmental Research* 121, 52–63.
- Longhurst, J.W.S., Irwin, J.G., Chatterton, T.J., Hayes, E.T., Leksmono, N.S., Symons, J.K., 2009. The development of effects-based air quality management regimes. *Atmospheric Environment* 43, 64–78.
- Longhurst, J.W.S., Lindley, S.J., Watson, A.F.R., Conlan, D.E., 1996. The introduction of local air quality management in the United Kingdom: A review and theoretical framework. *Atmospheric Environment* 30, 3975–3985.
- Luhar, A.K., Patil, R.S., 1989. A General Finite Line Source Model for vehicular pollution prediction. *Atmospheric Environment* 23, 555–562.
- Masih, A., Saini, R., Singhvi, R., Taneja, A., 2010. Concentrations, sources, and exposure profiles of polycyclic aromatic hydrocarbons (PAHs) in particulate matter (PM₁₀) in the north central part of India. *Environmental Monitoring and Assessment* 163, 421–431.
- Mayer, H., 1999. Air pollution in cities. *Atmospheric Environment* 33, 4029–4037.
- Mazzera, D.M., Lowenthal, D.H., Chow, J.C., Watson, J.G., 2001. Sources of PM₁₀ and sulfate aerosol at McMurdo Station, Antarctica. *Chemosphere* 45, 347–356.
- Menon-Choudhary, D., Shukla, P.R., 2009. An integrated strategy for urban Air Quality Management in India. *International Journal of Environment and Pollution* 39, 233–252.
- MEP (Ministry of Environment Protection), 2013. The State Council Issues Action Plan on Prevention and Control of Air Pollution Introducing Ten Measures to Improve Air Quality, http://english.mep.gov.cn/News_service/infocus/201309/t20130924_260707.htm, accessed in September 2014.
- Miller, C.A., Hidy, G., Hales, J., Kolb, C.E., Werner, A.S., Haneke, B., Parrish, D., Frey, H.C., Rojas-Bracho, L., Deslauriers, M., Pennell, B., Mobley, J.D., 2006. Air emission inventories in North America: A critical assessment. *Journal of the Air & Waste Management Association* 56, 1115–1129.
- MoEF (Ministry of Environment and Forest), 2009. *National Ambient Air Quality Standards*, Government of India, New Delhi, 4 pages.
- Mofarrah, A., Husain, T., 2010. A holistic approach for optimal design of air quality monitoring network expansion in an urban area. *Atmospheric Environment* 44, 432–440.
- Mohan, M., Kandy, A., 2007. An analysis of the annual and seasonal trends of air quality index of Delhi. *Environmental Monitoring and Assessment* 131, 267–277.
- Molina, L.T., Madronich, S., Gaffney, J.S., Apel, E., de Foy, B., Fast, J., Ferrare, R., Herndon, S., Jimenez, J.L., Lamb, B., Osornio-Vargas, A.R., Russell, P., Schauer, J.J., Stevens, P.S., Volkamer, R., Zavala, M., 2010. An overview of the MILAGRO 2006 Campaign: Mexico City emissions and their transport and transformation. *Atmospheric Chemistry and Physics* 10, 8697–8760.
- Molina, L.T., de Foy, B., Martinez, O.V., Paramo Figueroa, V.H., 2009. Air Quality, Weather and Climate in Mexico City, World Meteorological Organization (WMO) Bulletin 58, 48 pages.
- Molina, L.T., Kolb, C.E., de Foy, B., Brune, W.H., Jimenez, J.L., Ramos-Villegas, R., Sarmiento, J., Paramo-Figueroa, V.H., Cardenas, B., Gutierrez-Avedoy, V., Molina, M.J., 2007. Air quality in North America's most populous city & overview of the MCMA-2003 campaign. *Atmospheric Chemistry and Physics* 7, 2447–2473.
- Molina, L.T., Molina, M.J., 2002. *Air Quality in the Mexico Megacity: An Integrated Assessment*, Kluwer Academic Publishers, Dordrecht, The Netherlands, pp.141–145.
- Moorcroft, S., Dore, C., 2013. Review of Effectiveness of Local Authority Action Plans and Future Policy Options for LAQM, Prepared by Air Quality Consultant Ltd., Bristol UK on behalf of DEFRA, 64 pages.
- Moussiopoulos, N., 2003. *Air Quality in Cities*, Heidelberg, Springer, 299 pages.
- Moussiopoulos, N., Kalognomou, E.A., Douros, I., Samaras, Z., Giannouli, M., Mellios, G., 2005. Air pollution level at hotspot areas of selected European cities. *Proceeding of 10th Conference on Harmonization within Atmospheric Dispersion Modelling for Regulatory Purposes*, Sissi, Crete, pp. 283–287.

- Mustafic, H., Jabre, P., Caussin, C., Murad, M.H., Escolano, S., Tafflet, M., Perier, M.C., Marijon, E., Vernerey, D., Empana, J.P., Jouven, X., 2012. Main air pollutants and myocardial infarction: a systematic review and meta-analysis. *JAMA—Journal of the American Medical Association* 307, 713–721.
- Nagendra, S., Khare, M., Gulia, S., Vijay, P., Chithra, V.S., Bell, M., Namdeo, A., 2012. Application of ADMS and AERMOD models to study the dispersion of vehicular pollutants in urban areas of India and the United Kingdom. *Proceeding of 20th WIT International Conference on Air Pollution*, May 16–18, 2012, Coruna, Spain, pp. 3–12.
- Nagpure, A.S., Gurjar, B.R., 2012. Development and evaluation of Vehicular Air Pollution Inventory model. *Atmospheric Environment* 59, 160–169.
- Naiker, Y., Diab, R.D., Zunckel, M., Hayes, E.T., 2012. Introduction of local air quality management in South Africa: Overview and challenges. *Environmental Science & Policy* 17, 62–71.
- NAAQs-USA (National ambient air quality standards for USA), 2014. <http://www.epa.gov/air/criteria.html>, accessed in March 2014.
- NAQO-UK (National air quality objectives for UK), 2014. http://uk-air.defra.gov.uk/documents/National_air_quality_objectives.pdf, accessed in April 2014.
- NEA (National Environment Agency), 2014. <http://app2.nea.gov.sg/anti-pollution-radiation-protection/air-pollution-control>, accessed in March 2014.
- Newcastle City Council, 2006. Air Quality Action Plan of Newcastle City Centre AQMA, Prepared by Air Quality Consultant, UK and University of West of England, Bristol, UK, pp. 48 pages.
- NILU (Norwegian Institute for Air Research), 2007. Air Quality Management Plan for e-Thekwini Municipality, Kwa-Zulu Natal, South Africa, Produced by e-Thekwini Health and Norwegian Institute for Air Research, 79 pages.
- Norman, R., Cairncross, E., Witi, J., Bradshaw, D., 2007. Estimating the burden of disease attributable to urban outdoor air pollution in South Africa in 2000. *SAMJ South African Medical Journal* 97, 782–790.
- NRC (National Research Council), 2004. Executive summary of air quality management in the United States, committee on *Air Quality Management in the United States, Executive summary*, 32 pages.
- NSW (New South Wales) Government, 2010. Current Air Quality in New South Wales, Australia – A Technical Paper Supporting the Clean Air Forum 2010, DECCW 2010/728, 58 pages.
- NSW (New South Wales) Government, 2009. Action for Air, Department of Environment, Climate Change and Water NSW, Report no. 712, Australia, 32 pages.
- Oliveira, C., Pio, C., Caseiro, A., Santos, P., Nunes, T., Mao, H.J., Luahana, L., Sokhi, R., 2010. Road traffic impact on urban atmospheric aerosol loading at Oporto, Portugal. *Atmospheric Environment* 44, 3147–3158.
- Pakbin, P., Ning, Z., Shafer, M.M., Schauer, J.J., Sioutas, C., 2011. Seasonal and spatial coarse particle elemental concentrations in the Los Angeles area. *Aerosol Science and Technology* 45, 1528–1529.
- Pandolfi, M., Gonzalez-Castanedo, Y., Alastuey, A., de la Rosa, J.D., Mantilla, E., de la Campa, A.S., Querol, X., Pey, J., Amato, F., Moreno, T., 2011. Source apportionment of PM₁₀ and PM_{2.5} at multiple sites in the strait of Gibraltar by PMF: Impact of shipping emissions. *Environmental Science and Pollution Research* 18, 260–269.
- Pant, P., Harrison, R.M., 2012. Critical review of receptor modelling for particulate matter: A case study of India. *Atmospheric Environment* 49, 1–12.
- Pant, P., Harrison, R.M., 2013. Estimation of the contribution of road traffic emissions to particulate matter concentrations from field measurements: A review. *Atmospheric Environment* 77, 78–97.
- Parrish, D.D., Singh, H.B., Molina, L., Madronich, S., 2011. Air quality progress in North American megacities: A review. *Atmospheric Environment* 45, 7015–7025.
- Pascal, M., Corso, M., Chanel, O., Declercq, C., Badaloni, C., Cesaroni, G., Henschel, S., Meister, K., Haluza, D., Martin-Ormedo, P., Medina, S., Grp, A., 2013. Assessing the public health impacts of urban air pollution in 25 European cities: Results of the Aphekom project. *Science of the Total Environment* 449, 390–400.
- Patil, R.S., Kumar, R., Menon, R., Shah, M.K., Sethi, V., 2013. Development of particulate matter speciation profiles for major sources in six cities in India. *Atmospheric Research* 132, 1–11.
- Peel, J.L., Klein, M., Flanders, W.D., Mulholland, J.A., Tolbert, P.E., 2010. Impact of improved air quality during the 1996 Summer Olympic Games in Atlanta on multiple cardiovascular and respiratory outcomes. *Research Report- Health Effect Institute* 148, 3–23.
- Pererz, L., Medina-Ramon, M., Kunzli, N., Alastuey, A., Pey, J., Perez, N., Garcia, R., Tobias, A., Querol, X., Sunyer, J., 2009. Size fractionated particulate matter, vehicle traffic, and case-specific daily mortality in Barcelona, Spain. *Environmental Science & Technology* 43, 4707–4714.
- Perrone, M.G., Larsen, B.R., Ferrero, L., Sangiorgi, G., De Gennaro, G., Udisti, R., Zangrando, R., Gambaro, A., Bolzacchini, E., 2012. Sources of high PM_{2.5} concentrations in Milan, Northern Italy: Molecular marker data and CMB modelling. *Science of the Total Environment* 414, 343–355.
- Pope, C.A., Dockery, D.W., 2006. Health effects of fine particulate air pollution: Lines that connect. *Journal of the Air & Waste Management Association* 56, 709–742.
- Pope, C.A., Ezzati, M., Dockery, D.W., 2009. Fine-particulate air pollution and life expectancy in the United States. *New England Journal of Medicine* 360, 376–386.
- Preuthippan, A., Udomsubpayakul, U., Chaisupamongkollarp, T.C., Pentamwa, P., 2004. Effect of PM₁₀ pollution in Bangkok on children with and without asthma. *Pediatric Pulmonology* 37, 187–192.
- Pundir, B.P., Das, S., 1985. A State of the Art Report on Vehicle Emissions, Department of Environment, Government of India, Project No. K-12011/3/83/EN 1, New Delhi.
- Qin, Y.J., Kim, E., Hopke, P.K., 2006. The concentrations and sources of PM_{2.5} in metropolitan New York City. *Atmospheric Environment* 40, S312–S332.
- Raaschou-Nielsen, O., Andersen, Z.J., Beelen, R., Samoli, E., Stafoggia, M., Weinmayr, G., Hoffmann, B., Fischer, P., Nieuwenhuijsen, M.J., Brunekreef, B., Xun, W.W., Katsouyanni, K., Dimakopoulou, K., Sommar, J., Forsberg, B., Modig, L., Oudin, A., Oftedal, B., Schwarze, P.E., Nafstad, P., De Faire, U., Pedersen, N.L., Ostenson, C.G., Fratiglioni, L., Penell, J., Korek, M., Pershagen, G., Eriksen, K.T., Sorensen, M., Tjonneland, A., Ellermann, T., Eeftens, M., Peeters, P.H., Meliefste, K., Wang, M., Bueno-de-Mesquita, B., Key, T.J., de Hoogh, K., Concin, H., Nagel, G., Vilier, A., Gironi, S., Krogh, V., Tsai, M.Y., Ricceri, F., Sacerdote, C., Galassi, C., Migliore, E., Ranzi, A., Cesaroni, G., Badaloni, C., Forastiere, F., Tamayo, I., Amiano, P., Dorronsoro, M., Trichopoulos, A., Bamia, C., Vineis, P., Hoek, G., 2013. Air pollution and lung cancer incidence in 17 European cohorts: Prospective analyses from the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Lancet Oncology* 14, 813–822.
- Ramachandra, T.V., Shwetmala, 2009. Emissions from India's transport sector: Statewise synthesis. *Atmospheric Environment* 43, 5510–5517.
- Ramadan, Z., Eickhout, B., Song, X.H., Buydens, L.M.C., Hopke, P.K., 2003. Comparison of Positive Matrix Factorization and Multilinear Engine for the source apportionment of particulate pollutants. *Chemometrics and Intelligent Laboratory Systems* 66, 15–28.
- Salcedo, D., Castro, T., Ruiz-Suarez, L.G., Garcia-Reynoso, A., Torres-Jardon, R., Torres-Jaramillo, A., Mar-Morales, B.E., Salcido, A., Celada, A.T., Carreon-Sierra, S., Martinez, A.P., Fentanes-Arriaga, O.A., Deustua, E., Ramos-Villegas, R., Retama-Hernandez, A., Saavedra, M.I., Suarez-Lastra, M., 2012. Study of the regional air quality south of Mexico City (Morelos state). *Science of the Total Environment* 414, 417–432.
- Santosa, S.J., Okuda, T., Tanaka, S., 2008. Air pollution and urban air quality management in Indonesia. *Clean-Soil Air Water* 36, 466–475.
- Scire, J.S., Strimaitis D.G., Yamartino, R.J., 2000. A user's Guide for the CALPUFF Dispersion Model (Version 5), Earth Tech Inc., Miami, United States, 552 pages.

- Schauer, J.J., Cass, G.R., 2000. Source apportionment of wintertime gas-phase and particle-phase air pollutants using organic compounds as tracers. *Environmental Science & Technology* 34, 1821–1832.
- Schauer, J.J., Rogge, W.F., Hildemann, L.M., Mazurek, M.A., Cass, G.R., Simoneit, B.R.T., 1996. Source apportionment of airborne particulate matter using organic compounds as tracers. *Atmospheric Environment* 30, 3837–3855.
- Schipa, I., Mangia, C., Tanzarella, A., Conte, D., Marra, G.P., Rizza, U., 2011. A GIS based air quality system for the Apulia region, southern Italy. *International Journal of Environment and Pollution* 47, 120–133.
- Scorgie, Y., 2012. *Urban Air Quality Management and Planning in South Africa*. Ph.D. Thesis, University of Johannesburg, Department of Geography, Environmental Management and Energy Studies, Johannesburg, South Africa, 333 pages.
- SEI (Stockholm Environment Institute), 2004. A Strategic Framework for Air Quality Management in Asia, Part of Air Pollution in the Megacities of Asia (APMA) Project and the Clean Air Initiative for Asian Cities (CAI-Asia), 45 pages.
- Sharma, S.K., Mandal, T.K., Saxena, M., Sharma, R.A., Datta, A., Saud, T., 2014. Variation of OC, EC, WSIC and trace metals of PM₁₀ in Delhi, India. *Journal of Atmospheric and Solar–Terrestrial Physics* 113, 10–22.
- Sharma, N., Gulia, S., Dhyani, R., Singh, A., 2013a. Performance evaluation of CALINE 4 dispersion model for an urban highway corridor in Delhi. *Journal of Scientific & Industrial Research* 72, 521–530.
- Sharma, S., Sharma, P., Khare, M., 2013b. Hybrid modelling approach for effective simulation of reactive pollutants like Ozone. *Atmospheric Environment* 80, 408–414.
- Sharma, H., Jain, V.K., Khan, Z.H., 2007. Characterization and source identification of polycyclic aromatic hydrocarbons (PAHs) in the urban environment of Delhi. *Chemosphere* 66, 302–310.
- Sharma, D.N., Sawant, A.A., Uma, R., Cocker, D.R., 2003. Preliminary chemical characterization of particle-phase organic compounds in New Delhi, India. *Atmospheric Environment* 37, 4317–4323.
- Shasha, D., 2010. http://news.xinhuanet.com/english2010/china/2010-11/25/c_13622601.htm, accessed in November 2012.
- Shridhar, V., Khillare, P.S., Agarwal, T., Ray, S., 2010. Metallic species in ambient particulate matter at rural and urban location of Delhi. *Journal of Hazardous Materials* 175, 600–607.
- Simoneit, B.R.T., Schauer, J.J., Nolte, C.G., Oros, D.R., Elias, V.O., Fraser, M.P., Rogge, W.F., Cass, G.R., 1999. Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles. *Atmospheric Environment* 33, 173–182.
- Simpson, R., Williams, G., Petroeschovsky, A., Best, T., Morgan, G., Denison, L., Hinwood, A., Neville, G., Neller, A., 2005. The short-term effects of air pollution on daily mortality in four Australian cities. *Australian and New Zealand Journal of Public Health* 29, 205–212.
- Singh, A.K., Gupta, H.K., Gupta, K., Singh, P., Gupta, V.B., Sharma, R.C., 2007. A comparative study of air pollution in Indian cities. *Bulletin of Environmental Contamination and Toxicology* 78, 411–416.
- Sivertsen, B., 2008. Monitoring air quality, objectives and design. *Chemical Industry & Chemical Engineering Quarterly* 14, 167–171.
- Sivertsen, B., Bartonova, A., 2012. Air Quality Management Planning (AQMP). *Chemical Industry & Chemical Engineering Quarterly* 18, 667–674.
- Sonawane, N.V., Patil, R.S., Sethi, V., 2012. Health benefit modelling and optimization of vehicular pollution control strategies. *Atmospheric Environment* 60, 193–201.
- Song, Y., Zhang, Y.H., Xie, S.D., Zeng, L.M., Zheng, M., Salmon, L.G., Shao, M., Slanina, S., 2006. Source apportionment of PM_{2.5} in Beijing by positive matrix factorization. *Atmospheric Environment* 40, 1526–1537.
- Soret, A., Jimenez-Guerrero, P., Andres, D., Cardenas, F., Rueda, S., Baldasano, J.M., 2013. Estimation of future emission scenarios for analysing the impact of traffic mobility on a large Mediterranean conurbation in the Barcelona Metropolitan Area (Spain). *Atmospheric Pollution Research* 4, 22–32.
- Soret, A., Jimenez-Guerrero, P., Baldasano, J.M., 2011. Comprehensive air quality planning for the Barcelona Metropolitan Area through traffic management. *Atmospheric Pollution Research* 2, 255–266.
- Srimuruganandam, B., Nagendra, S.M.S., 2011. Chemical characterization of PM₁₀ and PM_{2.5} mass concentrations emitted by heterogeneous traffic. *Science of the Total Environment* 409, 3144–3157.
- Srivastava, A., Jain, V.K., 2008. Source apportionment of suspended particulate matters in a clean area of Delhi: A note. *Transportation Research Part D—Transport and Environment* 13, 59–63.
- Srivastava, A., Gupta, S., Jain, V.K., 2008. Source apportionment of total suspended particulate matter in coarse and fine size ranges over Delhi. *Aerosol and Air Quality Research* 8, 188–200.
- Stankovic, S., Vaskovic, V., Petrovic, N., Radojicic, Z., 2012. Sustainable air pollution management in urban areas caused by traffic: Case study Banja Luka. *Technics Technologies Education Management—TTEM* 7, 1615–1619.
- Steinar, L., Grqnskei, K.E., Hannegraaf, M.C., Jansen, H., Kuik, O.J., Oosterhuis, F.H., Olsthoorn, X.A., Jitendra, J.S., Tanvi, N., Carter, J.B., 1997. *Urban Air Quality Management Strategy in Asia – Guidebook*, The World Bank Press, Washington D.C., pp. 192.
- Stone, E., Schauer, J., Quraishi, T.A., Mahmood, A., 2010. Chemical characterization and source apportionment of fine and coarse particulate matter in Lahore, Pakistan. *Atmospheric Environment* 44, 1062–1070.
- Sturtz, T.M., Adar, S.D., Gould, T., Larson, T.V., 2014. Constrained source apportionment of coarse particulate matter and selected trace elements in three cities from the multi-ethnic study of atherosclerosis. *Atmospheric Environment* 84, 65–77.
- Tao, J., Cheng, T.T., Zhang, R.J., Cao, J.J., Zhu, L.H., Wang, Q.Y., Luo, L., Zhang, L.M., 2013. Chemical composition of PM_{2.5} at an urban site of Chengdu in southwestern China. *Advances in Atmospheric Sciences* 30, 1070–1084.
- Tecer, L.H., Tuncel, G., Karaca, F., Alagha, O., Suren, P., Zararsiz, A., Kirmaz, R., 2012. Metallic composition and source apportionment of fine and coarse particles using positive matrix factorization in the Southern Black Sea atmosphere. *Atmospheric Research* 118, 153–169.
- Terblanche, A.P., Opperman, L., Nel, C.M., Reinach, S.G., Tosen, G., Cadman, A., 1992. Preliminary results of exposure measurements and health effects of the vaal triangle air pollution health study. *South African Medical Journal* 81, 550–556.
- The Hindu, 2014. <http://www.thehindu.com/sci-tech/health/only-12-percent-of-citydwellers-breathe-safe-air-says-who/article5986549.ece>, accessed in May 2014.
- Thurston, G.D., Ito, K., Lall, R., 2011. A source apportionment of U.S. fine particulate matter air pollution. *Atmospheric Environment* 45, 3924–3936.
- Thurston, G.D., Ozkaynak, H., Schatz, A., 1984. A chemical characterization and source apportionment of the IP Network fine particle data. *Proceeding of the 77th Annual Meeting of the Air Pollution Control Association*, June 24–29, 1984, San Francisco, CA, United States.
- Tie, X.X., Madronich, S., Li, G.H., Ying, Z.M., Zhang, R.Y., Garcia, A.R., Lee-Taylor, J., Liu, Y.B., 2007. Characterizations of chemical oxidants in Mexico city: A regional chemical dynamical model (WRF–Chem) study. *Atmospheric Environment* 41, 1989–2008.
- Tiwari, S., Srivastava, A.K., Bisht, D.S., Safai, P.D., Parmita, P., 2013. Assessment of carbonaceous aerosol over Delhi in the Indo-Gangetic Basin: characterization, sources and temporal variability. *Natural Hazards* 65, 1745–1764.
- Todd, J., Dodds, J., Goetzmann, M., Jennings, P., Scally, G., Lukatelich, R., Graham-Taylor, S., Smith, R., Upton, M., Heymans, A., Olsen, B., 2011. Implementing the Perth Air Quality Management Plan, 6th Progress Report of the Air Quality Control Committee, 108 pages.
- TOI (Times of India), 2014. India’s Air Quality Among Five Worst, Daily Newspaper, The Times of India, 30th January 2014, New Delhi.

- Tonne, C., Beevers, S., Kelly, F.J., Jarup, L., Wilkinson, P., Armstrong, B., 2010. An approach for estimating the health effects of changes over time in air pollution: An illustration using cardio-respiratory hospital admissions in London. *Occupational and Environmental Medicine* 67, 422–427.
- Tonne, C., Beevers, S., Armstrong, B., Kelly, F., Wilkinson, P., 2008. Air pollution and mortality benefits of the London Congestion Charge: Spatial and socioeconomic inequalities. *Occupational and Environmental Medicine* 65, 620–627.
- UN–Habitat, 2010. The State of Asian Cities 2010/11, United Nations Human Settlements Programme, HS/162/10 E, Japan, 280 pages.
- U.S. Department of Energy, 2011. Reducing Petroleum Dependence in On-Road Transportation in the United States, Clean Cities Five-Year Strategic Plan, Department of Energy, United States, 47 pages.
- U.S. EPA (U.S. Environment Protection Agency), 2012. Our Nation's Air Status and Trends through 2010, EPA-454/R-12-001, 32 pages.
- U.S. EPA (U.S. Environment Protection Agency), 2011a. http://www.epa.gov/oaqps001/aqportal/management/aq_goalsetting.htm. Accessed in September 2011.
- U.S. EPA (U.S. Environment Protection Agency), 2011b. <http://www.epa.gov/otaq/models.htm>, accessed in November 2011.
- U.S. EPA (U.S. Environment Protection Agency), 2011c. http://www.epa.gov/air/aqportal/management/public_participation.htm, accessed in November 2011.
- U.S. EPA (U.S. Environment Protection Agency), 2005. Federal Register, Part –III, 40 CFR Part 51, Revisions to the Guideline on Air Quality Models: Adoption of a Preferred General Purpose (Flat and Complex Terrain) Dispersion Model and Other Revisions; Final Rule, 45 pages.
- U.S. EPA (U.S. Environmental Protection Agency), 2002. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=29060>, accessed in September 2014.
- Viana, M., Kuhlbusch, T.A.J., Querol, X., Alastuey, A., Harrison, R.M., Hopke, P.K., Winiwarter, W., Vallius, A., Szidat, S., Prevot, A.S.H., Hueglin, C., Bloemen, H., Wahlin, P., Vecchi, R., Miranda, A.I., Kasper-Giebl, A., Maenhaut, W., Hitenberger, R., 2008. Source apportionment of particulate matter in Europe: A review of methods and results. *Journal of Aerosol Science* 39, 827–849.
- Vlachokostas, C., Achillas, C., Moussiopoulos, N., Hourdakos, E., Tsiligiridis, G., Ntziachristos, L., Baniyas, G., Stavrakakis, N., Sidiropoulos, C., 2009. Decision support system for the evaluation of urban air pollution control options: Application for particulate pollution in Thessaloniki, Greece. *Science of the Total Environment* 407, 5937–5948.
- Wang, S.X., Hao, J.M., 2012. Air quality management in China: Issues, challenges, and options. *Journal of Environmental Sciences–China* 24, 2–13.
- Wang, Z.B., Hu, M., Wu, Z.J., Yue, D.L., He, L.Y., Huang, X.F., Liu, X.G., Wiedensohler, A., 2013. Long-term measurements of particle number size distributions and the relationships with air mass history and source apportionment in the summer of Beijing. *Atmospheric Chemistry and Physics* 13, 10159–10170.
- Wang, H., Fu, L., Zhou, Y., Du, X., Ge, W., 2010. Trends in vehicular emissions in China's mega cities from 1995 to 2005. *Environmental Pollution* 158, 394–400.
- Watson, J.G., Chen, L.W.A., Chow, J.C., Doraiswamy, P., Lowenthal, D.H., 2008. Source apportionment: Findings from the U.S. Supersites program. *Journal of the Air & Waste Management Association* 58, 265–288.
- Watson, J.G., Zhu, T., Chow, J.C., Engelbrecht, J., Fujita, E.M., Wilson, W.E., 2002. Receptor modeling application framework for particle source apportionment. *Chemosphere* 49, 1093–1136.
- Watson, J.G., Chow, J.C., Houck, J.E., 2001. PM_{2.5} chemical source profiles for vehicle exhaust, vegetative burning, geological material, and coal burning in Northwestern Colorado during 1995. *Chemosphere* 43, 1141–1151.
- Wichmann, J., Voyi, K.V.V., 2005. Air pollution epidemiological studies in South Africa: Need for freshening up. *Epidemiology* 16, S20–S20.
- Williams, M., Barrowcliffe, R., Laxen, D., Monks, P., 2011. Report on Review of Air Quality Modelling in DEFRA by Air Quality Modelling Review Steering Group, King's College, London, 44 pages.
- Williams, R.J., Bhattacharyya, K.K., 2004. Provincial Framework for Airshed Planning in British Columbia, Ministry of Water, Land and Air Protection, Government of British Columbia, 43 pages.
- Wilson, R.D., 1997. Guidance on Incorporating Voluntary Mobile Source Emission Reduction Programs in State Implementation Plan, U.S. Environment Protection Agency, 15 pages.
- WHO (World Health Organization), 2014. <http://www.who.int/mediacentre/news/releases/2014/air-pollution/en/>, accessed in September 2014.
- WHO (World Health Organization), 2005. WHO Air Quality Guidelines for Particulate Matter, Ozone, Nitrogen Dioxide and Sulphur Dioxide, Global update 2005, World Health Organization, Geneva, 22 pages.
- Woodfield, N.K., Longhurst, J.W.S., Beattie, C.I., Chatterton, T., Laxen, D.P.H., 2006. Regional collaborative urban air quality management: Case studies across Great Britain. *Environmental Modelling & Software* 21, 595–599.
- Woodfield, N.K., Longhurst, J.W.S., Beattie, C.I., Laxen, D.P.H., 2003. Critical evaluation of the role of scientific analysis in UK local authority AQMA decision-making: Method development and preliminary results. *Science of the Total Environment* 311, 1–18.
- Wolf, M.E., Fields, P.G., Manne, G.K., Villegas, M.T.L., Bravo, V.G., Gomez, R.I., Coyoacan, D., 2009. Developing Mexico national emissions inventory projections for the future years of 2008, 2012, and 2030. *18th Annual International Emission Inventory Conference*, U.S. Environmental Protection Agency, Baltimore, USA.
- Wu, Y., Wang, R.J., Zhou, Y., Lin, B.H., Fu, L.X., He, K.B., Hao, J.M., 2011. On-road vehicle emission control in Beijing: Past, present, and future. *Environmental Science & Technology* 45, 147–153.
- Xie, S.D., Liu, Z., Chen, T., Hua, L., 2008. Spatiotemporal variations of ambient PM₁₀ source contributions in Beijing in 2004 using positive matrix factorization. *Atmospheric Chemistry and Physics* 8, 2701–2716.
- Yin, J.X., Harrison, R.M., Chen, Q., Rutter, A., Schauer, J.J., 2010. Source apportionment of fine particles at urban background and rural sites in the UK atmosphere. *Atmospheric Environment* 44, 841–851.
- Ying, Z.M., Tie, X.X., Li, G.H., 2009. Sensitivity of ozone concentrations to diurnal variations of surface emissions in Mexico City: A WRF/Chem modeling study. *Atmospheric Environment* 43, 851–859.
- Zheng, M., Salmon, L.G., Schauer, J.J., Zeng, L.M., Kiang, C.S., Zhang, Y.H., Cass, G.R., 2005. Seasonal trends in PM_{2.5} source contributions in Beijing, China. *Atmospheric Environment* 39, 3967–3976.
- Zhong, L.J., Louie, P.K.K., Zheng, J.Y., Wai, K.M., Ho, J.W.K., Yuan, Z.B., Lau, A.K.H., Yue, D.L., Zhou, Y., 2013a. The Pearl River Delta Regional Air Quality Monitoring Network – Regional Collaborative Efforts on Joint Air Quality Management. *Aerosol and Air Quality Research* 13, 1582–1582.
- Zhong, L.J., Louie, P.K.K., Zheng, J.Y., Yuan, Z.B., Yue, D.L., Ho, J.W.K., Lau, A.K.H., 2013b. Science policy interplay: Air quality management in the Pearl River Delta region and Hong Kong. *Atmospheric Environment* 76, 3–10.
- Zhou, Y., Wu, Y., Yang, L., Fu, L.X., He, K.B., Wang, S.X., Hao, J.M., Chen, J.C., Li, C.Y., 2010. The impact of transportation control measures on emission reductions during the 2008 Olympic Games in Beijing, China. *Atmospheric Environment* 44, 285–293.
- Zhu, L., Huang, X., Shi, H., Cai, X., Song, Y., 2011. Transport pathways and potential sources of PM₁₀ in Beijing. *Atmospheric Environment* 45, 594–604.
- Zunckel, M., 2009. Scoping Report on Air Quality Modelling Activities in Southern Africa and the Feasibility of a Regional Modelling Centre, Air Pollution Information Network – Africa, Natural Resources and the Environment, Congella, South Africa, 27 pages.